

INFINITE-DIMENSIONAL UNIFORM POLYHEDRA

SERGEY A. MELIKHOV

ABSTRACT. Uniform covers with a finite-dimensional nerve are rare (i.e., do not form a cofinal family) in many separable metric spaces of interest. To get hold on uniform homotopy properties of these spaces, a reasonably behaved notion of an infinite-dimensional metric polyhedron is needed; a specific list of desired properties was sketched by J. R. Isbell in a series of publications in 1959-64. In this paper we construct what appears to be the desired theory of uniform polyhedra; incidentally, considerable information about their metric and Lipschitz properties is obtained.

1. INTRODUCTION

The purpose of this paper is to develop a theory of polyhedra in the uniform category, thereby fulfilling what seems to be the basic aims of a research program initiated by J. R. Isbell in the 1950s. Specific terms of his program will be reviewed at the end of this introduction. This paper is a common sequel to [12; Chapter 2] and [13] and will in turn be followed by applications to (i) the Hilbert–Smith conjecture; (ii) $\varprojlim \varinjlim$ versus $\varinjlim \varprojlim$ of groups and Borsuk’s problem on embeddability of compact absolute retracts; (iii) Sklyarenko’s problem on shape invariance of Steenrod–Sitnikov homology.

1.1. What is going on in this paper. Since topological and uniform notions agree on compact spaces (recall that continuous maps with compact domain are uniformly continuous), the theory of uniform polyhedra is not supposed to say anything new about compact polyhedra. Moreover, there is nothing deep about finite-dimensional uniform polyhedra: any reasonable attempt to define them is expected to succeed, and end up with just an equivalent form of what Isbell himself once did in a few pages [8].

A key difficulty in the infinite-dimensional case can be seen from the following example. Let Δ^n be the standard n -simplex in $\mathbb{R}^{n+1}$, that is the intersection of the first octant with the hyperplane $\sum x_i = 1$. Then Δ^n has a constant (i.e. independent of n) edge length in Euclidean, or l_1 , or l_∞ metric. However, the distance from the barycenter of Δ^n (at $(\frac{1}{n+1}, \dots, \frac{1}{n+1})$) to the barycenter of a facet of Δ^n (at $(0, \frac{1}{n}, \dots, \frac{1}{n})$) tends to zero as $n \rightarrow \infty$, in either metric.

Thus barycentric subdivision does not work uniformly. It seems to be a difficult geometric problem whether there exists *any* subdivision of simplices into simplices with the property that for some metric (say l_1 , but not necessarily linear in general), the cover

Supported by Russian Foundation for Basic Research Grant No. 11-01-00822, Russian Government project 11.G34.31.0053 and Federal Program “Scientific and scientific-pedagogical staff of innovative Russia” 2009–2013.

by the open stars of vertices in the n th iterate of the subdivision refines the cover C_{ε_n} by all balls of radius ε_n , yet is refined by C_{δ_n} , where $\delta_n > 0$ and $\varepsilon_n \rightarrow 0$ as $n \rightarrow \infty$.

Our basic approach is to take a detour and use the “canonical subdivision” of [12], which when applied to a simplicial complex produces a cubical complex (versions of this construction are well-known in Geometric Group Theory and in Topological Combinatorics); each cube is then endowed with the l_∞ metric. This yields a combinatorially controlled uniform structure, since when applied to a cubical complex, the canonical subdivision is the obvious silly procedure producing another cubical complex by cutting every n -cube into 2^n of n -cubes. This is how we arrive at the uniform geometric realization of a simplicial complex.

While this simple idea already suffices for many practical purposes, it also brings some deep combinatorial complications. At least at a first glance, the “cubical” uniform structure appears to be in an endless conflict with basic PL constructions such as cone, join and mapping cylinder, which are manifestly “non-cubical”. It is this conflict that in a sense is the main subject of the present paper.

1.2. Why the heck is it all in terms of posets. Our first step towards resolution of the conflict is to further subdivide the cubes into simplices, without introducing new vertices. These simplices are now asymmetric, but each comes with a natural total order on its vertices, for it is isometric to the “standard skew n -simplex” $\{(x_1, \dots, x_n) \mid 0 \leq x_1 \leq \dots \leq x_n \leq 1\} \subset \mathbb{R}^n$ for some n with the l_∞ metric. One can now use these asymmetric metric simplices as separate building blocks, making sure their vertex orderings agree whenever they overlap. Thus let us call a partial ordering of the vertices of a simplicial complex *compatible* if it induces a total ordering on each simplex, or equivalently if vertices connected by an edge are comparable.

If K is a simplicial complex with a compatible partial ordering of vertices, or more generally any Δ -set (=semi-simplicial set in a modern terminology), then we can canonically endow every simplex of K with the standard skew metric; as long as there are no loops in the 1-skeleton (in the case of a Δ -set), this can be shown to extend to a path metric on the entire K . Unfortunately, this metric is generally plagued by the very same problem that we intended to avoid: it turns out that for each $\varepsilon > 0$ there exists an n such that every point of the standard skew n -simplex is ε -close to some point in its boundary (see Example 4.11).

For this reason we restrict our attention to flag complexes (a simplicial complex K is called “flag” if every subcomplex of K isomorphic to the boundary of a d -simplex, $d > 1$, lies in an actual d -simplex in K). A flag simplicial complex with a compatible partial ordering of vertices will be called a *preposet*. Note that the 1-skeleton of a preposet carries the structure of an acyclic digraph (=a directed graph with no directed cycles). Conversely, the flag complex spanned by an acyclic digraph is a preposet, with vertices partially ordered by the relation: $v \leq w$ if and only if there exists a directed path from v to w in the 1-skeleton.

A special case of a preposet is (the order complex of) a poset. Thus we arrive at the uniform geometric realization of a poset P . If P happens to be the poset of all nonempty faces of a simplicial complex K , this simply brings us back to the uniform geometric realization of K , as discussed above.

1.A. Main results

All posets and other combinatorial objects will be assumed countable throughout.

We consider three notions of uniform geometric realization of a poset (in particular, of the poset of nonempty faces of a simplicial or cubical complex) by a separable metrizable uniform space:

- by constructing an explicit embedding into the unit cube of the functional space c_0 (generalizing a construction of Shtan'ko–Shtogrin [14]);
- by gluing together the standard skew simplices via quotient uniformity (akin to the traditional geometric realization of a simplicial set via quotient topology);
- by gluing together the standard skew simplices via path metric (like in geometric polyhedral complexes used in Geometric Group Theory).

The following follows from Theorems 2.6 and 3.1:

Theorem 1.3. *All three notions of geometric realization are equivalent.*

The geometric realization of a locally infinite dimensional poset (even a simplicial complex) may fail to be complete; however it is “homotopy complete” in the sense that the completion can be instantaneously taken off the remainder by a homotopy (Lemma 4.7).

Geometric realization is promoted to a functor from monotone maps between posets to uniformly continuous maps between separable metrizable uniform spaces, which is shown to preserve pullbacks and those pushouts that remain pushouts upon barycentric subdivision (Theorem 3.8). In particular, the functor respects joins, and mapping cylinders of simplicial maps. Here the join of posets can refer to any of the two well-known distinct notions, and the join and mapping cylinder of metrizable uniform spaces are defined in [13].

To include arbitrary pushouts, such as mapping cylinders of general monotone maps, the geometric realization functor has to be extended to preposets. Let us emphasize that the mapping cylinder fails to be a poset (and so is only a preposet) for some monotone maps between posets that arise naturally in practice:

- the diagonal embedding $P \rightarrow P \times P$ (see [12; Example 2.14]);
- approximations to uniformly continuous maps (see Theorem 7.4);
- bonding maps between nerves of covers (see Example 6.19 and Lemma 8.2).

One can, however, emulate the mapping cylinder of a monotone map between posets by a certain poset, without changing the relative uniform homotopy equivalence class of the geometric realization (Corollary 4.15).

Unfortunately, preposets do not quite live up to our expectations. It turns out that there exists a preposet X whose geometric realization is not uniformly locally contractible; worse yet, it contains essential loops of arbitrarily small diameters (Example 5.2). In fact, $|X|$ is not uniformly homotopy equivalent to the geometric realization of any poset (Theorem 5.3).

Theorem 1.4. *The geometric realization of every poset P is uniformly locally contractible.*

This is saying that for each $\varepsilon > 0$ there exists a $\delta > 0$ such that every two δ -close uniformly continuous maps from an arbitrary metric space into $|P|$ are uniformly ε -homotopic with values in $|P|$ (Theorem 5.1).

On the other hand, there exists a poset Y whose geometric realization is not a uniform ANR; worse yet, for each $\varepsilon > 0$ it contains an embedded sphere of some dimension that is essential in $|Y|$, but null-homotopic in the ε -neighborhood of $|Y|$ in some fixed uniform ANR (Example 7.5). In fact, $|Y|$ is not uniformly homotopy equivalent to the geometric realization of any *conditionally complete* poset (Theorem 7.6), that is, a poset where every nonempty set has either the least upper bound or no upper bound whatsoever.

Conditionally complete posets include (the posets of nonempty faces of) simplicial and cubical complexes but exclude for instance (those of) some simplicial pseudo-complexes in the sense of Hilton–Wylie (where each simplex is embedded, but different simplices may have more than one face in common). Fortunately, the geometric realization of a conditionally complete poset does turn out to be a uniform ANR (Corollary 7.2). It is the geometric realizations of conditionally complete posets that we call *uniform polyhedra*.

Theorem 1.5. *Uniform polyhedra are uniform ANRs.*

This is saying that whenever a uniform polyhedron is uniformly embedded as a closed subset of a metrizable uniform space X , then it is a uniform retract of some its uniform neighborhood in X .¹

Theorem 1.5 is arguably the hardest result of the paper. The case of simplicial complexes is somewhat easier (Theorem 4.10), and actually reduces to the case of “cubohedra”, which was already treated in [13; Theorem 4.28]. (A cubohedron is a cubical complex that is a subcomplex of the standard cubical lattice in c_0 .) The finite-dimensional simplicial case is equivalent to a result of Isbell [8; Theorem 1.9], taking into account that his geometric realization is uniformly homeomorphic to ours in the case of a finite-dimensional simplicial complex.

Simplicial and/or cubical complexes alone do not form a closed theory with respect to a sufficient supply of operations that work uniformly. The mapping cylinder of a simplicial map, while being a poset, is certainly not a simplicial complex in general. Worse yet, it need not even be a conditionally complete poset ([12; Example 2.68]). This is arguably the single most important deficiency of our theory; but up to uniform

¹An examination of the proof reveals that uniform polyhedra endowed with the path metric are in fact Lipschitz ANRs. It seems unlikely that they can be remetrized as 1-Lipschitz ANRs.

homotopy, there is a remarkable workaround. The *thickened mapping cylinder* $TMC(f)$ of a monotone map $f: P \rightarrow Q$ between posets is a natural subset of the join $P * Q$, related to the graph of f (see [12]). If P and Q are conditionally complete, then so is $TMC(f)$, and up to relative uniform homotopy equivalence of geometric realizations it is the same as the usual mapping cylinder (Theorem 4.12). Using this, we obtain the following result (Theorem 8.10):

Theorem 1.6. *If X is a uniform ANR, then $X \times \mathbb{R}$ is uniformly homotopy equivalent to a uniform polyhedron.*

We also establish what appears to be the ultimate inverse limit representation theorem, and a uniform analogue of Hanner’s characterization of ANRs (Theorems 8.1 and 8.7):

Theorem 1.7. (a) *Every separable metrizable complete uniform space is the limit of an inverse sequence of uniformly continuous maps between geometric realizations of simplicial complexes.*

(b) *A separable metrizable uniform space is a uniform ANR if and only if it is uniformly ε -homotopy dominated by the geometric realization of a simplicial complex for each $\varepsilon > 0$.*

Weaker forms of (a) and (b), with cubohedra in place of simplicial complexes, are contained in the author’s previous paper [13; Corollary 4.30 and Theorem 5.13]. The present versions, being based on nerves of uniform covers rather than uniform neighborhoods in c_0 , have the advantage of greater flexibility, so they can be straightforwardly adapted, for instance, to equivariant contexts.

The compact case of (a) is due to Freudenthal [7], though the basic idea goes back to Alexandroff’s earlier work (see Remark 8.4). The residually finite-dimensional case of (a) is equivalent to a result of Isbell [10; Lemma V.33] (see also [4; Lemma 1.6], [11; Lemma 14]), though he made a minor mistake in his proof, which can be corrected as we show in the proof of Theorem 8.1. The residually finite-dimensional case of the “only if” direction in (b) is equivalent to another result of Isbell [9; 7.3].

1.B. Isbell’s problem

Let us now discuss how our results address Isbell’s research program. One attempted formulation appears in his book “Uniform spaces”:

“Research Problem B_2 . INFINITE-DIMENSIONAL POLYHEDRA. There is a large problem here, namely the systematic investigation of topological and uniform realizations of abstract simplicial complexes. One important paper in the literature (Dowker [1952]) has examined this problem, not from a categorical viewpoint. Dowker’s work tends to confirm, what many successful applications suggest, that for topology J. H. C. Whitehead’s realization by CW-complexes has strong claims to preference. Its definition is as simple as could be: [...] But Dowker’s work highlights the point that the suitability of CW-complexes for homology and homotopy is not conclusive; many realizations are topologically distinct but homotopy equivalent.

By now substantial experience in uniform spaces supports the pretensions of uniform complexes, in the finite-dimensional case only. (In any case they are homotopy equivalent (topologically) with CW-complexes; Dowker [1952].) In general they are not satisfactory, e.g. because they lack subdivisions. One can save the subdivisions, or any sufficiently narrow requirement, by tailoring a definition to fit. (Kuzminov and Švedov [1960] define a realization for which IV.6 [the covers by the stars of vertices in iterated barycentric subdivisions form a basis of the uniformity] is always valid; but all their applications are in the finite-dimensional case.) The real problem holding up progress is, what applications can be made of infinite-dimensional polyhedra in the general theory of uniform spaces? It would probably be beside the point to carry out a formal investigation of realizations with no specific applications in mind.” [10] (1964)

Comments: (i) As stated, the problem is quite vague, but some clarification on what kind of infinite-dimensional uniform complexes are sought here can be inferred from Isbell’s previous comments in his earlier papers, quoted below.

(ii) The covers by the stars of vertices in iterated *canonical* subdivisions do form a basis of the uniformity of our uniform polyhedra (see Theorem 3.2).

“It should be noted that the theorem [that Isbell’s finite-dimensional uniform simplicial complexes are complete uniform ANRs] as stated is trivially false for arbitrary uniform complexes, since some of them are incomplete. It is false for many complete ones also. It seems likely that strong results might be gotten by using some suitable uniformity for a complex, different from the one defined by $\max |x_\alpha - y_\alpha|$, though not necessarily different for finite-dimensional complexes.” [8] (1959)

Comments: indeed, with our adjusted uniformity, the theorem is now extended to infinite-dimensional simplicial complexes (Theorem 4.10), apart from the completeness.

Isbell only considered complete uniform ANRs, as well as their non-metrizable generalization, called ANRUs. Indeed he showed that completeness is forced by including certain non-metrizable spaces in the setup. However, if one works in the category of metrizable uniform spaces, then there is no such restriction, as noticed independently by G. L. Garg and N. T. Nhu in the 1970s (see [13; §4.B]). The resulting theory of (possibly non-complete) uniform ANRs has been developed only recently but turns out to be very flexible [13]. These uniform ANRs are homotopy complete (i.e. the completion can be instantaneously taken off the remainder) [13; Theorem 4.9], which for all practical purposes makes them just as easily manageable as if they were complete.

“I should like to repeat the remark from [[4] and [8]] that the uniform complexes are clearly not the right concept for the infinite-dimensional case. The finite-dimensionality in 7.2 [that every residually finite-dimensional complete uniform space X is an inverse limit (i) of finite-dimensional uniform simplicial complexes, (ii) of inverse limits of nerves of appropriate uniform covers of X] and 7.3 [that every residually finite-dimensional ANRU is controlled uniformly homotopy dominated by a finite dimensional uniform simplicial complex] may very likely appear for no better reason than that we do not have the right uniformity for the complexes.” [9] (1961)

Comments: indeed, the said results extend to all separable uniform spaces, using our adjusted uniformity on (countable) infinite-dimensional simplicial complexes (Corollary 8.5 and Theorems 8.6 and 8.8), except that cubical, rather than simplicial, complexes are needed for the generalization of 7.2(i).

Neither of the results extends to inseparable uniform spaces by the negative solution to Isbell’s “Research Problem B_3 ”, obtained independently by J. Pelant and E. V. Shchepin in 1975, and clarified recently by A. Hohti (see references in [13; §2.E]). Hohti’s result is that the unit ball of the inseparable space l_∞ is not point-finite, i.e. does not have a basis of uniform covers that have a finite multiplicity at each point.

1.C. Acknowledgments

I would like to thank P. M. Akhmetiev, N. Brodsky, A. N. Dranishnikov, J. Dydak, S. Illman, E. V. Shchepin and M. Skopenkov for useful discussions.

In what follows we assume some degree of familiarity with [12; Chapter 2] and the entire paper [13].

2. GEOMETRIC REALIZATION VIA EMBEDDING

Rectilinear geometric realization of a *finite* preposet is described in [12; §1.C]. [12; Lemma 2.11] yields a realization of an arbitrary preposet within some *combinatorial* simplex. This however avoids the issue of sensible geometric realization of the simplex itself. Let us now address it. In the case of finite simplicial complexes, the following construction yields essentially the same result as in [14]. (The author found it while being unaware of [14].)

2.1. Geometric realization. If S is a set, the functional space $[0, 1]^S$ of all maps $f: S \rightarrow [0, 1]$ is endowed with the metric $d(f, g) = \sup_{s \in S} |f(s) - g(s)|$. Note that the underlying uniform space of $[0, 1]^S$ is just $U(S, [0, 1])$, where S endowed with the discrete uniformity. The subset $\{0, 1\}^S$ of $[0, 1]^S$ may be identified with the power set 2^S , by associating to every subset $T \subset S$ its *characteristic function* $\chi_T: S \rightarrow [0, 1]$, defined by $\chi_T(T) = 1$ and $\chi_T(S \setminus T) = 0$. Note that if S is finite, $[0, 1]^S$ is just the usual $|S|$ -dimensional cube with the l_∞ metric, and $\{0, 1\}^S$ is the set of its vertices.

Let us recall the embedding of a poset into a simplex given by [12; Lemma 2.11(b)]. Given a poset $P = (\mathcal{P}, \leq)$, we identify every $p \in P$ with the cone $[p]$, viewed as an element of $2^{\mathcal{P}} = \{0, 1\}^{\mathcal{P}} \subset [0, 1]^{\mathcal{P}}$. Then the *geometric realization* of P is a subspace

$|P| \subset [0, 1]^{\mathcal{P}}$, defined to be the union of the convex hulls of all nonempty finite chains of P . Note that the cube vertex at the origin, $\{0, 0, \dots\}$ is never in $|P|$ since $[p]$ is never empty.

More generally, given a preposet $P = (\mathcal{P}, \prec)$, following [12; Lemma 2.11(a)] we inject it into its transitive closure $\langle P \rangle = (\mathcal{P}, \prec\prec)$, and identify every $p \in P$ with the cone $[p]_{\prec\prec}$ in the transitive closure, viewed again as an element of $2^{\mathcal{P}} = \{0, 1\}^{\mathcal{P}} \subset [0, 1]^{\mathcal{P}}$. Then the *geometric realization* of P is a subspace $|P| \subset [0, 1]^{\mathcal{P}}$, defined to be the union of the convex hulls of all nonempty finite chains of P . Every chain of P is a chain of $\langle P \rangle$, hence $|P| \subset |\langle P \rangle|$.

Since $[0, 1]^{\mathcal{P}}$ is complete, the closure $\overline{|P|}$ of $|P|$ in $[0, 1]^{\mathcal{P}}$ is uniformly homeomorphic to the completion of $|P|$. Note that each convex hull in the definition of $|P|$ is compact, and therefore separable. Hence if P is countable, $|P|$ is separable; consequently $\overline{|P|}$ is a Polish uniform space, that is, a separable metrizable complete uniform space.

2.2. Generalized geometric realization. Let P be a preposet, and fix an injection $j: P \rightarrow 2^S$ for some S . The underlying set 2^S of the poset 2^S is identified, as before, with $\{0, 1\}^S \subset [0, 1]^S$. Then the *geometric j -realization* of P is a subspace $|P|_j \subset [0, 1]^S$, defined to be the union of the convex hulls of the j -images of all nonempty finite chains in P . We note four basic examples:

- The injection j_P of [12; Lemma 2.11] yields the *standard* geometric realization $|P|_{j_P} = |P|$.
- If P has a least element, then there is a more economical embedding $j'_P: P \hookrightarrow 2^{\partial^* P}$, $p \mapsto [p] \cap \partial^* P$. We call $|P|' := |P|_{j'_P}$ the *reduced* geometric realization of P .
- On the other hand, if P is an atomic poset, then [12; Lemma 2.25] yields a more economical embedding $a_P: P \hookrightarrow \Delta^{A(P)}$. We call $|P|^\bullet := |P|_{a_P}$ the *atomic* geometric realization of P . Note that $a_P: \Delta^\Lambda \rightarrow \Delta^\Lambda$ is the identity.
- Finally if $P = C^*Q$, where Q is an atomic poset, then a_Q extends to $a'_P: P \hookrightarrow 2^{A(P)}$. We still call $|P|^\bullet := |P|_{a'_P}$ the *atomic* geometric realization of P . Note that $a'_{2^\Lambda}: 2^\Lambda \rightarrow 2^\Lambda$ is the identity.

2.3. Geometric realization of cone precomplex. If $P = (\mathcal{P}, \leq)$ is a cone complex, then $\mathcal{P}$ is countable, and the cone $[p]$ of every $p \in P$ is finite. Then $j_P(P) \subset \Delta^{\mathcal{P}}$ lies in the weak $\mathcal{P}$ -simplex $\Delta_w^{\mathcal{P}}$, that is the set of all nonempty finite subsets of $\mathcal{P}$ (see [12; 2.53]). If C is a finite chain of $\Delta^{\mathcal{P}}$, its convex hull lies in $[0, 1]^{\sup C} \times \{0\}^{\mathcal{P} \setminus \sup C}$. Consequently, $|\Delta_w^{\mathcal{P}}|^\bullet$ lies in $q_0 := ([0, 1], 0)^{(\mathcal{P}^+, \infty)}$, where $\mathcal{P}^+ = \mathcal{P} \cup \{\infty\}$ is the one-point compactification of the discrete space $\mathcal{P}$. Since q_0 is complete, $\overline{|\Delta_w^{\mathcal{P}}|^\bullet}$ also lies in q_0 . More generally, if P is a cone precomplex, its transitive closure $\langle P \rangle$ is a cone complex, and $\overline{|P|}$ lies in $\overline{|\langle P \rangle|} \subset \overline{|\Delta_w^{\mathcal{P}}|^\bullet} \subset q_0$.

It is not hard to see that q_0 itself is identified with $\overline{|2_{\mathcal{P}}^{\mathcal{P}}|^\bullet}$. On the other hand, note that $\overline{|\Delta_{\mathbb{N}}^{\mathbb{N}}|^\bullet} \subsetneq \overline{|\Delta^{\mathbb{N}}|^\bullet}$, since $|\Delta^{\mathbb{N}}|^\bullet$ includes convex hulls of finite chains of infinite subsets of $\mathbb{N}$. In fact, since there are uncountably many of infinite subsets of $\mathbb{N}$, $|\Delta^{\mathbb{N}}|^\bullet$ is not separable.

If P is a cone complex that is *Noetherian*, i.e. contains no infinite chain (which could only be ascending since all cones are finite), then $|P|$ is closed in $[0, 1]^P$ and hence complete. It follows that $|P|$ is complete also for every Noetherian cone *precomplex* P , that is a preposet whose transitive closure is a Noetherian cone complex.

Example 2.4. If P is a poset and $j: P \rightarrow 2^S$ is an injection but not an embedding, then $|P|_j$ need not be isometric to $|P|$. Indeed, let P be the subposet of $2^{\{a,b,c\}}$ with elements $\emptyset, \{a\}, \{a, b\}, \{a, b, c\}, \{c\}$. Let $j: P \rightarrow 2^{\{a,b,c\}}$ re-embed $\{c\}$ onto $\{b\}$ and fix the other elements. Let C be the chain $\{\emptyset, \{a\}, \{a, b\}, \{a, b, c\}\}$ and let D be the chain $\{\emptyset, \{c\}, \{a, b, c\}\}$ in P . A point $x \in |C| \subset |P|$ has coordinates (x_a, x_b, x_c) for some numbers $1 \geq x_a \geq x_b \geq x_c \geq 0$, and a point $y' \in |D|$ has coordinates (y_1, y_2) for some numbers $1 \geq y_2 \geq y_1 \geq 0$. Then the image y of y' in $|P|$ has coordinates (y_2, y_2, y_1) , and the image y_j of y' in $|P|_j$ has coordinates (y_2, y_1, y_2) . Setting $(x_a, x_b, x_c) = (\frac{3}{4}, \frac{1}{2}, \frac{1}{4})$ and $(y_1, y_2) = (\frac{3}{4}, \frac{1}{2})$, we obtain $d(x, y) = \frac{1}{2}$ and $d(x, y_j) = \frac{1}{4}$.

Example 2.5. Here is a simpler example of the same kind. Let P be the subposet of $2^{\{a,b\}}$ with elements $\emptyset, \{a\}$ and $\{b\}$. Let $j: P \rightarrow 2^{\{a,b\}}$ re-embed $\{a\}$ onto $\{a, b\}$ and fix the other elements. Let C be the chain $\{\emptyset, \{a\}\}$ and let D be the singleton chain $\{\{b\}\}$ in P . Let $x \in |C| \subset |P|$ have coordinates $(x_a, x_b) = (0, \frac{1}{2})$; then its image x_j in $|P|_j$ has coordinates $(\frac{1}{2}, \frac{1}{2})$. The point $y \in |D| = \{y\}$ has coordinates $(1, 0)$. Hence $d(x, y) = 1$ and $d(x_j, y) = \frac{1}{2}$.

Theorem 2.6. *If P is a poset and $j: P \rightarrow 2^S$ is an embedding, then $|P|_j$ is isometric to $|P|$.*

This trivially implies that if P is a preposet and $j: P \rightarrow 2^S$ is an injection that factors through an embedding of the transitive closure, then $|P|_j$ is isometric to $|P|$.

Proof. We first consider the case where P is the totally ordered n -element poset $[n] = (\{1, \dots, n\}, \leq)$, where $\leq$ has the usual meaning. To avoid confusion, we consider the standard embedding $j_{[n]}$ of $[n]$ in $2^{\{1, \dots, n\}}$, that is, $j_{[n]}(i) = \{1, \dots, i\}$. Then each $j_{[n]}(i) \in 2^{\{1, \dots, n\}}$ is identified with the vertex $(0, \dots, 0, 1, \dots, 1)$ ($n-i$ zeroes, i ones) of the simplex $||[n]|| \subset [0, 1]^n$. Similarly each $P_i := j(i) \in 2^S$ is identified with a point of $\{0, 1\}^S \subset [0, 1]^S$. Then $j j_{[n]}^{-1}$ extends uniquely to an affine map $\Phi_j: ||[n]|| \rightarrow ||[n]||_j$. A point $x = (x_1, \dots, x_n) \in [0, 1]^n$ lies in $||[n]||$ if and only if $1 = x_1 \geq x_2 \geq \dots \geq x_n \geq 0$. Let us write $\hat{x} = \Phi_j(x)$. It is easy to see that $\hat{x}(s) = x_i$ for each $s \in P_i \setminus P_{i-1}$, where $P_0 = \emptyset$, and $\hat{x}(s) = 0$ for $s \notin P_n$.

Given another point $y \in ||[n]||$, since each $P_i \setminus P_{i-1}$ is nonempty, it follows that $d(\hat{x}, \hat{y}) = \sup_{s \in S} |\hat{x}(s) - \hat{y}(s)|$ equals $\max_{i \in [n]} |x_i - y_i| = d(x, y)$. Thus Φ_j is an isometry, which completes the proof of the case $P = [n]$.

We now resume the proof of the general case, where $P = (\mathcal{P}, \leq)$ is an arbitrary poset. Given a finite chain C in P , we may represent C as the image of the poset $[k]$, where k is the cardinality of C , under the (unique) isomorphism $c: [k] \rightarrow C$. Consider the compositions $c: [k] \xrightarrow{c} P \subset 2^P$ and $j c: [k] \xrightarrow{c} P \xrightarrow{j} 2^S$. These extend to the

isometries $\Phi_c: |[k]| \rightarrow |[k]|_c \subset |P|$ and $\Phi_{jc}: |[k]| \rightarrow |[k]|_{jc} \subset |P|_j$. The compositions $|[k]|_c \xrightarrow{\Phi_c^{-1}} |[k]| \xrightarrow{\Phi_{jc}} |[k]|_{jc}$ agree with each other for different c , and thus combine into a map $\Phi_j: |P| \rightarrow |P|_j$ that is an isometry on the convex hull of every finite chain of P .

To complete the proof, it suffices to show that all $x, y \in |P|$ satisfy $d(\Phi_j(x), \Phi_j(y)) = d(x, y)$. This will follow once we prove that $d(\Phi_j(x), \Phi_j(y))$ does not depend on j . We may assume without loss of generality that P has the least element $\hat{0}$ and the greatest element $\hat{1}$; for if P has no least (resp. greatest) element, then $S \notin j(P)$ (resp. $\emptyset \notin j(P)$), and therefore j extends to an embedding of CP (resp. C^*P) in 2^S defined by $\hat{1} \mapsto S$ (resp. $\hat{0} \mapsto \emptyset$). Let A (resp. B) be some chain in P containing $\hat{0}$ and $\hat{1}$, whose convex hull contains x (resp. y). We consider the (unique) isomorphisms $a: [m] \rightarrow A$ and $b: [n] \rightarrow B$, where m is the cardinality of A and n is the cardinality of B . Thus $a(1) = \hat{0} = b(1)$ and $a(m) = \hat{1} = b(n)$.

Let $\prec$ be the covering relation of the subposet $A \cup B$ of P . (That is, $x, y \in A \cup B$ satisfy $x \prec y$ iff $x < y$ and there exists no $z \in A \cup B$ such that $x < z < y$.) Let $(k_1, l_1), \dots, (k_r, l_r)$ be all pairs in $[m] \times [n]$ such that either $a(k_i) = b(l_i)$ or $a(k_i) \notin B$, $b(l_i) \notin A$ and $a(k_i) \prec b(l_i)$, where each $k_i \leq k_{i+1}$. Let $Z \subset [r]$ be the set of indices i such that $a(k_i) = b(l_i)$. It is easy to see² that each $k_i < k_{i+1}$ and each $l_i < l_{i+1}$. Similarly let $(k'_1, l'_1), \dots, (k'_{r'}, l'_{r'})$ be all pairs in $[m] \times [n]$ such that either $a(k'_i) = b(l'_i)$ or $a(k'_i) \notin B$, $b(l'_i) \notin A$ and $a(k'_i) \succ b(l'_i)$; we may assume that each $k'_i < k'_{i+1}$ and each $l'_i < l'_{i+1}$. Let Z' be the set of indices i such that $a(k'_i) = b(l'_i)$. We note that $k_1 = l_1 = k'_1 = l'_1 = 1$, $k_r = k'_{r'} = m$ and $l_r = l'_{r'} = n$.

It is easy to see³ that $a(k_i + 1) \not\leq b(l_{i+1} - 1)$ for each i . Let us write $A_i = ja(i)$ and $B_i = jb(i)$. Since j is an embedding, we obtain that $A_{k_i+1} \not\subset B_{l_{i+1}-1}$. On the other hand, since $k_i < k_{i+1}$, we have $A_{k_i+1} \subset A_{k_{i+1}} \subset B_{l_{i+1}}$; and similarly $A_{k_i} \subset B_{l_{i+1}-1}$. Thus $A_{k_i+1} \setminus A_{k_i}$ has a nonempty intersection with $B_{l_{i+1}} \setminus B_{l_{i+1}-1}$. In other words, the set Σ of all pairs (κ, λ) such that $A_\kappa \setminus A_{\kappa-1}$ has a nonempty intersection with $B_\lambda \setminus B_{\lambda-1}$ includes the set Δ of all pairs of the form $(k_i + 1, l_{i+1})$. By symmetry, Σ also includes the set Δ' of all pairs of the form $(k'_{i+1}, l'_i + 1)$.

We claim that for each $(\kappa, \lambda) \in \Sigma$ there exist a $(k, l) \in \Delta$ and a $(k', l') \in \Delta'$ such that $k \leq \kappa \leq k'$ and $l' \leq \lambda \leq l$. Indeed, if $a(\kappa) \leq b(\lambda - 1)$, then $A_\kappa \subset B_{\lambda-1}$; in particular, $(\kappa, \lambda) \notin \Sigma$. If $a(\kappa)$ and $b(\lambda - 1)$ are incomparable, let i be the maximal number such that

²Indeed, suppose that $k_i = k_{i+1}$ (and $l_i \neq l_{i+1}$). The cases (1) $i, i+1 \in Z$; (2) $i \in Z$ and $i+1 \notin Z$; (3) $i \notin Z$ and $i+1 \in Z$ are ruled out for trivial reasons. In the remaining case (4) $i, i+1 \notin Z$ we have either $l_i < l_{i+1}$ or $l_i > l_{i+1}$. Then either $a(k_{i+1}) = a(k_i) < b(l_i) < b(l_{i+1})$ or $a(k_i) = a(k_{i+1}) < b(l_{i+1}) < b(l_i)$. Hence either $a(k_{i+1}) \not\leq b(l_{i+1})$ or $a(k_i) \not\leq b(l_i)$, which is a contradiction. Thus $k_i < k_{i+1}$. Next suppose that $l_i \geq l_{i+1}$. Then $a(k_i) < a(k_{i+1}) \leq b(l_{i+1}) \leq b(l_i)$, so $i \notin Z$. Hence $a(k_i) < a(k_{i+1}) < b(l_{i+1}) \leq b(l_i)$, so $a(k_i) \not\leq b(l_i)$, which is a contradiction.

³Indeed, if $a(k_i + 1) = b(l_{i+1} - 1)$, then $k_i + 1 = k_j$ and $l_{i+1} - 1 = l_j$ for some j ; hence $i < j < i + 1$, which is a contradiction. Suppose that $a(k_i + 1) < b(l_{i+1} - 1)$. Since $<$ is the transitive closure of $\prec$, there exist $\kappa \geq k_i + 1$ and $\lambda \leq l_{i+1} - 1$ such that $a(\kappa) \prec b(\lambda)$. If $a(\kappa) \in B$, then $k_i < \kappa = k_j$ for some j such that $b(l_j) = a(k_j) = a(\kappa) < b(\lambda)$. Hence $l_j < \lambda < l_{i+1}$, and therefore $i < j < i + 1$, which is a contradiction. Thus $a(\kappa) \notin B$, and similarly $b(\lambda) \notin A$. Hence $k_i < \kappa = k_j$ and $l_{i+1} > \lambda = l_j$ for some j ; hence $i < j < i + 1$, which is a contradiction.

$\kappa > k_i$. Then $\kappa \leq k_{i+1}$, so $\lambda - 1 < l_{i+1}$. Finally, if $a(\kappa) > b(\lambda - 1)$, let i be the maximal number such that $\kappa > k_i$. We claim that still $\lambda - 1 < l_{i+1}$. Suppose on the contrary that $\lambda - 1 \geq l_{i+1}$. Then $a(\kappa) > b(\lambda - 1) \geq b(l_{i+1}) \geq a(k_{i+1})$. On the other hand, $\kappa \leq k_{i+1}$ by our choice of i , and so $a(\kappa) \leq a(k_{i+1})$, which is a contradiction. This completes the proof of the assertion on (k, l) ; and the assertion on (k', l') is proved similarly.

We have $x = \Phi_a(\alpha)$ and $y = \Phi_b(\beta)$ for some $\alpha = (\alpha_1, \dots, \alpha_m) \in |A|$ and some $\beta = (\beta_1, \dots, \beta_n) \in |B|$. Let us denote $\Phi_j(x) = \Phi_{ja}(\alpha)$ by $\hat{\alpha}$ and $\Phi_j(y) = \Phi_{jb}(\beta)$ by $\hat{\beta}$. We have $d(\hat{\alpha}, \hat{\beta}) = \sup_{s \in S} |\hat{\alpha}(s) - \hat{\beta}(s)|$. Here $\hat{\alpha}(s) = \alpha_\kappa$ for each $s \in A_\kappa \setminus A_{\kappa-1}$ (where $A_0 = \emptyset$) and $\hat{\alpha}(s) = 0$ for $s \notin A_m$; similarly, $\hat{\beta}(s) = \beta_\lambda$ for each $s \in B_\lambda \setminus B_{\lambda-1}$ (where $B_0 = \emptyset$) and $\hat{\beta}(s) = 0$ for $s \notin B_n$. Hence $d(\hat{\alpha}, \hat{\beta}) = \max_{(\kappa, \lambda) \in \Sigma} |\alpha_\kappa - \beta_\lambda|$.

Next we recall that $1 = \alpha_1 \geq \alpha_2 \geq \dots \geq \alpha_m \geq 0$ and $1 = \beta_1 \geq \beta_2 \geq \dots \geq \beta_n \geq 0$. In particular, $k \leq \kappa \leq k'$ implies $\alpha_k \geq \alpha_\kappa \geq \alpha_{k'}$; and $l' \leq \lambda \leq l$ implies $\beta_{l'} \geq \beta_\lambda \geq \beta_l$. Hence $\alpha_k - \beta_l \geq \alpha_\kappa - \beta_\lambda \geq \alpha_{k'} - \beta_{l'}$, which implies $|\alpha_\kappa - \beta_\lambda| \leq \max\{|\alpha_k - \beta_l|, |\alpha_{k'} - \beta_{l'}|\}$. Thus

$$d(\hat{\alpha}, \hat{\beta}) = \max_{(k, l) \in \Delta \cup \Delta'} |\alpha_k - \beta_l|.$$

The right hand side does not depend on j ; therefore so does the left hand side, that is, $d(\Phi_j(x), \Phi_j(y))$. $\square$

Corollary 2.7. *If Q is a subpreposet of a preposet P , then $|Q|$ admits a natural isometric embedding in $|P|$.*

Proof. If P and Q are posets, then by Theorem 2.6, $|Q| = |Q|_{j_Q}$ is isometric to $|Q|_j \subset |P|_{j_P} = |P|$, where j is the composition $Q \subset P \xrightarrow{j_P} 2^P$.

In the general case, the transitive closure $\langle Q \rangle$ is a subposet of $\langle P \rangle$, and it is easy to see that the image of the isometric embedding $|Q| \subset |\langle Q \rangle| \rightarrow |\langle P \rangle|$ lies in $|P|$. $\square$

Corollary 2.8. *If P is a preposet, $|P^*|$ is isometric to $|P|$.*

Proof. First assume that P is a poset. By Theorem 2.6, there exists an isometry $\Phi_j: |P| \rightarrow |P|_j$, where $j: P \rightarrow 2^P$ is defined by $j(p) = \mathcal{P} \setminus \{p\}$. The isomorphism $\varphi: 2^P \rightarrow (2^P)^*$ defined by $\varphi(S) = \mathcal{P} \setminus S$ extends to a self-isometry Φ of $[0, 1]^P$, taking $|P|_j$ onto $|P^*|$. Indeed we have $\varphi j(p) = j_{P^*}(p^*)$, where $j_{P^*}: P^* \rightarrow 2^P$ is the standard embedding, $j_{P^*}(p^*) = \lceil p^* \rceil = \lfloor p \rfloor^*$.

If P is a preposet, we apply the above construction to its transitive closure $\langle P \rangle$. Given a finite chain of P , viewed as an embedding $c: [n] \rightarrow \langle P \rangle \subset 2^P$, we have $\lfloor [n] \rfloor_c \subset |P| \subset |\langle P \rangle|$. Clearly, the isometry $|\langle P \rangle| \rightarrow |\langle P \rangle|_j \rightarrow |\langle P \rangle^*| = |\langle P^* \rangle|$ takes $\lfloor [n] \rfloor_c$ onto $\lfloor [n]^* \rfloor_{c^*}$. $\square$

Remark 2.9. We recall that 2_w^P is identified with a closed subposet of $(I_c^P)^*$ (see [12; 2.53]). The completed geometric realization $\overline{|I_c^P|}$ therefore contains an isometric copy of $\overline{|2_w^P|}$. The latter is in turn isometric to the completed *atomic* geometric realization $\overline{|2_w^P|}^\bullet = q_0$. This isometry extends to an isometry between $\overline{|I_c^P|}$ and $Q_0 := ([-1, 1], 0)^{(\mathcal{P}^+, \infty)}$.

Theorem 2.10. *Let P and Q be preposets.*

- (a) $|P \times Q|$ is uniformly homeomorphic to $|P| \times |Q|$.
 (b) $|P + Q|$ is uniformly homeomorphic to $|P| * |Q|$.
 In particular, $|CP|$ and $|C^*P|$ are uniformly homeomorphic to $C|P|$.
 (c) $|P * Q|$ is uniformly homeomorphic to $|P| * |Q|$.

Proof. (a). The injections $P \rightarrow 2^P$ and $Q \rightarrow 2^Q$ as in [12; Lemma 2.11] yield $|P| \subset [0, 1]^P$ and $|Q| \subset [0, 1]^Q$, where $P = (\mathcal{P}, \preceq)$ and $Q = (\mathcal{Q}, \leq)$. The injection $P \times Q \rightarrow 2^P \times 2^Q \simeq 2^{P \sqcup Q}$ yields $|P \times Q| \subset [0, 1]^{P \sqcup Q}$. Meanwhile, $|P| \times |Q|$ lies in $[0, 1]^P \times [0, 1]^Q$, which may be identified with $[0, 1]^{P \sqcup Q}$. To see that $|P \times Q| = |P| \times |Q|$ under this identification, it suffices to consider the case where P and Q are nonempty finite totally ordered sets. This case (and the more general case where P and Q are finite) follows using that a chain in $2^P \times 2^Q = 2^{P \sqcup Q}$ lies in $P \times Q$ if and only if it projects onto a chain in P and onto a chain in Q . $\square$

(b). Consider the injection $P + Q \rightarrow 2^{P \sqcup pt \sqcup Q}$ defined by $\sigma \mapsto \lceil \sigma \rceil$ if $\sigma \in P$, and by $\sigma \mapsto \lceil \sigma \rceil \cup pt$ if $\sigma \in Q$. This yields $|P + Q| \subset [0, 1]^{P \sqcup pt \sqcup Q}$, so that $|P|$ lies in $[0, 1]^P \times \{0\} \times \{0\}$ and $|Q|$ in $\{1\} \times \{1\} \times [0, 1]^Q$. It is easy to see that $|P + Q|$ is the union of $|P|$, $|Q|$ and all straight line segments with one endpoint in $|P|$ and another in $|Q|$. (Beware that these segments alone cover $|P + Q|$ only if both P and Q are nonempty.) Thus $|P + Q|$ is the independent rectilinear join of $|P|$ and $|Q|$, as defined in [13; §3.B]. Hence by [13; Theorem 3.38], $|P + Q|$ is uniformly homeomorphic to $|P| * |Q|$. $\square$

(c). From definition, $P * Q$ is the subpreposet $C^*P \times Q \cup P \times C^*Q$ of $C^*P \times C^*Q$. By the proof of part (b), $|C^*P|$ is the rectilinear cone $c|P|$, as defined in [13; §3.B]. Then by part (a), $|C^*P \times C^*Q|$ is uniformly homeomorphic to $c|P| \times c|Q|$; whereas $|P * Q|$ is uniformly homeomorphic to its subspace $c|P| \times |Q| \cup |P| \times c|Q|$. Write $|P| = X$ and $|Q| = Y$ for the sake of brevity. Then, Lemmas 3.37, 3.35 and 3.29 in [13] yield uniform homeomorphisms

$$cX \times Y \cup X \times cY \rightarrow cX \times Y \bigcup_{X \times Y} X \times cY \rightarrow CX \times Y \bigcup_{X \times Y} X \times CY \rightarrow X * Y,$$

where each of the amalgamated unions in the middle is defined as a pushout in the category of uniform spaces (and so is endowed with the quotient uniformity). $\square$

3. GEOMETRIC REALIZATION VIA QUOTIENT

Given a collection of preposets $P_\alpha = (\mathcal{P}_\alpha, \leq)$, their *disjoint union* $\bigsqcup_\alpha P_\alpha$ is their coproduct in the category of preposets; more explicitly, it is the preposet $(\bigsqcup_\alpha \mathcal{P}_\alpha, \preceq)$, where $p_\alpha \in P_\alpha$ and $p_\beta \in P_\beta$ satisfy $p_\alpha \preceq p_\beta$ iff $\alpha = \beta$ and $p_\alpha \leq p_\beta$. We note that disjoint union does not commute with geometric realization unless the index set is finite, because every infinite disjoint union of non-discrete uniform spaces is easily seen to be non-metrizable.

Theorem 3.1. *Let P be a poset, and let $P_\sqcup$ be the disjoint union of all nonempty finite chains of P . Let $\pi: P_\sqcup \rightarrow P$ be determined by the inclusions $C \subset P$, where $C \in P^\flat$. Then*

- (a) $|\pi|: |P_\sqcup| \rightarrow |P|$ is a quotient map (in the category of uniform spaces);
 (b) if d is the standard metric on $|P_\sqcup|$, then

$$d_\infty(x, y) = \inf_{n \in \mathbb{N}} \inf_{\substack{x_1, \dots, x_{n-1} \in |P| \\ (x_0 := x, x_n := y)}} \sum_{i=0}^{n-1} d(\pi^{-1}(x_i), \pi^{-1}(x_{i+1}))$$

is a metric on $|P|$.

Theorem 3.1(a) implies that $|P|$ is a quotient space of $|P_\sqcup|$ (in the category of uniform spaces). This is reminiscent of the definition of geometric realization of semi-simplicial sets, and of the well-known characterization of the topology of a CW-complex as the topology of a quotient (in the category of topological spaces!) of the disjoint union of its cells.

Theorem 3.1(b) is reminiscent of the definition of geometric polyhedral complexes used in metric geometry and in geometric group theory (see [2], [3]).

Proof. This is based on the technique of quotient maps of finite type (see [13; §3.A]) and on the proof of Theorem 2.6 above.

Write $q = |\pi|$, and let d stand for the usual metric on $|P_\sqcup|$ and on P . Clearly, q is surjective. Given $x, y \in |P|$, let $d_n(x, y) = \inf_{x=x_0, \dots, x_n=y} \sum d(q^{-1}(x_i), q^{-1}(x_{i+1}))$ and $d_\infty(x, y) = \inf_{n \in \mathbb{N}} d_n(x, y)$. It is easy to see that d_∞ is a pseudo-metric on $|P|$ (while each d_n need not satisfy the triangle axiom) and that the identity maps $(|P|, d_n) \xrightarrow{\text{id}} (|P|, d_\infty) \xrightarrow{\text{id}} (|P|, d)$ are uniformly continuous for each n . If $(|P|, d) \xrightarrow{\text{id}} (|P|, d_n)$ is uniformly continuous for some n , then on the one hand, d_∞ is uniformly equivalent to d , and on the other hand, by [13; Theorem 3.17(b)], d_∞ induces the quotient uniformity on $|P|$. Thus it suffices to show that $(|P|, d) \xrightarrow{\text{id}} (|P|, d_3)$ is uniformly continuous.

Suppose that $P = (\mathcal{P}, \leq)$, and let $\hat{P} := C^*CP$ (with additional elements $\hat{0}$ and $\hat{1}$). The standard geometric realization $|P| \subset 2^{\mathcal{P}}$ lies in the reduced geometric realization $|\hat{P}'| \subset 2^{\mathcal{P} \cup \{\hat{1}\}}$ (where $\hat{0}$ is identified with $\emptyset \in 2^{\mathcal{P} \cup \{\hat{1}\}}$).

Pick some $x, y \in |P|$, and let $A \subset P$ and $B \subset P$ be any chains whose convex hulls contain x and y respectively. We have unique isomorphisms $a: [m] \rightarrow \hat{A}$ and $b: [n] \rightarrow \hat{B}$, where $a(1) = \hat{0} = b(1)$ and $a(m) = \hat{1} = b(n)$. Let $(k_1, l_1), \dots, (k_r, l_r)$ and $(k'_1, l'_1), \dots, (k'_{r'}, l'_{r'})$, and Z and Z' be as in the proof of Theorem 2.6. We also recall the notation $\alpha_i = x(s)$ for some $s \in a(i) \setminus a(i-1)$, and $\beta_j = y(t)$ for some $t \in b(j) \setminus b(j-1)$, where $2 \leq i \leq m$ and $2 \leq j \leq n$. Observe that this does not depend on the choices of s and t .

We now define an ‘intermediary’ chain $C \subset P$, viewed as an isomorphism $c: [q] \rightarrow \hat{C}$. The inductive construction starts with $c(1) = \hat{0}$, and in the event that $c(i) = \hat{1}$ it terminates with $q = i$. Suppose that $c(i) = a(k)$ for some $k < m$; if $k \neq k_j$ for any j , then let $c(i+1) = a(k+1)$; if $k = k_j$ for some $j \notin Z$, then let $c(i+1) = b(l_j)$. Similarly, suppose that $c(i) = b(l)$ for some $l < n$; if $l \neq l'_j$ for any j , then let $c(i+1) = b(l+1)$; if $l = l'_j$ for some $j \notin Z'$, then let $c(i+1) = a(k'_j)$. Finally, if $c(i) = a(k) = b(l) < \hat{1}$, then we are free to set either $c(i+1) = a(k+1)$ or $c(i+1) = b(l+1)$.

Next we define an $x' \in |A'|_{a'}$ and a $y' \in |B'|_{b'}$, where $A' = A \cap C$ and $B' = B \cap C$, viewed as isomorphisms $a': [m'] \rightarrow \hat{A}'$ and $b': [n'] \rightarrow \hat{B}'$. Given an $s \in \mathcal{P}$, we have $s \in a'(m') = [\hat{1}] = \mathcal{P} \cup \{\hat{1}\}$, and $s \notin a'(1) = [\hat{0}] = \emptyset$. Hence $s \in a'(i) \setminus a'(i-1)$, where $2 \leq i \leq m'$. Pick some $h_i \in [m]$ so that $a'(i-1) \leq a(h_i-1)$ and $a(h_i) \leq a'(i)$. We must be more specific for $i=1$ and $i=m$, and we set $h_2 = 2$ (which is the least among all possible choices) and $h_{m'} = m$ (which is the greatest among all possible choices). Since $h_i \geq 2$, we may set $x'(s) = \alpha_{h_i}$. Let $\alpha'_i = x'(s)$ for any $s \in a'(i) \setminus a'(i-1)$, where $2 \leq i \leq m'$; clearly this does not depend on the choice of s . Thus $\alpha'_2 = \alpha_2$ and $\alpha'_{m'} = m$. Since $x \in |A|_a$, we have $\alpha_2 = 1$ and $\alpha_m = 0$. Therefore $\alpha'_2 = 1$ and $\alpha'_{m'} = 0$, whence $x' \in |A'|_{a'}$. We can similarly define a $y' \in |B'|_{b'}$ and consequently β'_j where $2 \leq j \leq n'$.

Let us estimate $d(x, x')$ from above. Suppose that $s \in a'(i) \setminus a'(i-1)$, where $2 \leq i \leq m'$. If both $a'(i)$ and $a'(i-1)$ belong to $\hat{C}$, then $x'(s) = x(s)$. Else we have $a'(i-1) = a(k_j)$ and $a'(i) = a(k'_{j'})$, where $1 \leq j < r$ and $1 < j' \leq r'$. Moreover, by the construction of C we have $l'_{j'-1} < l_j \leq l'_{j'} < l_{j+1}$. By definition, both $x(s)$ and $x'(s)$ belong to $[\alpha_{k_j+1}, \alpha_{k'_{j'}}]$. Since $l'_{j'-1} + 1 \leq l_{j+1}$, we have $\alpha_{k'_{j'}} - \alpha_{k_j+1} \leq \alpha_{k'_{j'}} - \beta_{l'_{j'-1}+1} + \beta_{l_{j+1}} - \alpha_{k_j+1}$. However $|\alpha_{k'_{j'}} - \beta_{l'_{j'-1}+1}| \leq d(x, y)$ and $|\beta_{l_{j+1}} - \alpha_{k_j+1}| \leq d(x, y)$ by the proof of Theorem 2.6. Thus $|x(s) - x'(s)| \leq 2d(x, y)$. We have proved that $d(x, x') \leq 2d(x, y)$.

We have $d_3(x, y) \leq d(x, x') + d(x', y') + d(y', y)$. By the above, $d(x, x') \leq 2d(x, y)$, and similarly $d(y', y) \leq 2d(x, y)$. By the triangle axiom, $d(x', y') \leq d(x', x) + d(x, y) + d(y, y') \leq 5d(x, y)$. Hence $d_3(x, y) \leq 9d(x, y)$. Thus $(|P|, d) \rightarrow (|P|, d_3)$ is uniformly continuous. $\square$

Corollary 3.2. *If P is a preposet, then $|P^\#|$ is uniformly homeomorphic to $|P|$ by a homeomorphism $h: |P^\#| \rightarrow |P|$.*

Moreover, if P is a poset, then $d_\infty(x, y) = 2d_\infty(h(x), h(y))$ for all $x, y \in |P^\#|$.

Proof. Consider the map of sets $f: P^\# \rightarrow |P| = |P|_{j_P}$ defined by sending an element $[\sigma, \sigma] \in P^\#$ into the vertex $[\{\sigma\}]_{j_P}$ of $|P|_{j_P}$, and an element $[\sigma, \tau] \in P^\#$ with $\sigma < \tau$ into the central point of the straight line segment connecting the vertices $[\{\sigma\}]_{j_P}$ and $[\{\tau\}]_{j_P}$. A finite chain C of $P^\#$ is of the form $[\sigma_1, \tau_1] \subseteq \cdots \subseteq [\sigma_n, \tau_n]$, where $\sigma_1 \leq \cdots \leq \sigma_n \leq \tau_n \leq \cdots \leq \tau_1$. (We recall that when P is a poset, $\subseteq$ is just the usual inclusion.) By collapsing all the equality signs in the latter string of inequalities we obtain a string of strict inequalities, which represents a chain $\tilde{C}$ of P (of length $\geq n$). Then $f(C) \subset |\tilde{C}|_{j_P}$; hence f extends by linearity to a map $h: |P^\#| \rightarrow |P|$.

It is not hard to see that h is a bijection (note that this is well-known in the case where P is a poset, cf. [12]). Indeed, for every chain D of P , $f^{-1}(|D|_{j_P})$ can be identified with $D^\#$. So the assertion reduces to the case $P = [n]$, which can be checked directly.

To show that h is a uniform homeomorphism we may assume that P is a poset by considering the transitive closure. Then it suffices to prove that $d_\infty(x, y) = 2d_\infty(h(x), h(y))$ for all $x, y \in |P^\#|$. By the definition of the d_∞ metrics (with respect to the usual metrics d on $P_\sqcup$ and $(P^\#)_\sqcup$; see the statement of Theorem 3.1(b)), it suffices to prove this when $x, y \in |D^\#|$ for some chain D of P . So the assertion again reduces to the case $P = [n]$, which can be checked directly. $\square$

Corollary 3.3. *Let P be a poset, and let $P_\square$ be the disjoint union of all intervals of P . Let $\rho: P_\square \rightarrow P$ be determined by the inclusions $Q \subset P$, where $Q \in P^\#$. Then $|\rho|: |P_\square| \rightarrow |P|$ is a quotient map (in the category of uniform spaces).*

Proof. Consider the commutative square

$$\begin{array}{ccc} (P_\square)_\square & \longrightarrow & P_\square \\ \pi' \downarrow & & \downarrow \rho \\ P_\square & \xrightarrow{\pi} & P. \end{array}$$

Here π' is trivially a quotient map, and π is a quotient map by Theorem 3.1(a). Hence $\pi\pi'$ is a quotient map, and therefore so is ρ . $\square$

Remark 3.4. Corollary 3.3 is, in a sense, easier than Theorem 3.1, for it can also be proved as follows. One first shows that $|r_X|$ (defined in §4 below) is uniformly continuous without using Corollary 3.3; this can be done by writing an explicit formula for $|r_X|$ in coordinates. Next one observes that $|r_X^\#|: |P^{\#\#}| \rightarrow |P^\#|$ takes any pair of sufficiently close points onto a pair of points contained in $|[q]|$ for the same interval $q \in P^\#$. It then remains to use [13; Theorem 3.17(b)] in the same way as it is used in the proof of Theorem 3.1.

Remark 3.5. Corollary 3.3 suffices to show that $h: |P^\#| \rightarrow |P|$, as defined in the proof of Corollary 3.2, is a uniform homeomorphism. Indeed, writing $P = (S, \leq)$, the usual metric on $|P|$ is induced from that on $|2^S|^\bullet$ via the usual embedding $P \hookrightarrow 2^S$, and similarly the usual metric on $|P^\#|$ is induced from that on $|(2^S)^\#|$ (see [12; Lemma 2.11] and Corollary 2.7). So it suffices to show that $h: |(2^S)^\#| \rightarrow |2^S|^\bullet$ is a uniform homeomorphism. Let $Q = (2^S)^\#$. By Corollary 3.3, $(|Q|, d)$ is uniformly homeomorphic to $(|Q|, d_\infty)$, where d denotes the usual metric on $|Q|$ and on $|Q_\square|$, and d_∞ is as in the statement of Theorem 3.1(b). On the other hand, it is easy to see that $(|Q|, d_\infty)$ is isometric to $(|2^S|^\bullet, 2d)$, where d is the usual l_∞ metric on the cube $|2^S|^\bullet$. Hence $(|Q|, d)$ is uniformly homeomorphic to $(|2^S|^\bullet, d)$.

Remark 3.6. If P is a simple poset (see [12; 2.54]), then it is easy to see that the usual metrics d on $|P_\square|$ and on $|P_\square|$ lead to the same d_∞ metric on $|P|$. So Theorem 3.1 for such P can be recovered from Corollary 3.3.

3.7. Geometric realization of a monotone map. Given a monotone map $f: P \rightarrow Q$ between posets, it extends uniquely to a map $|f|: |P| \rightarrow |Q|$ that is affine on every convex hull of a chain. In fact, $|f|$ is clearly 1-Lipschitz on every convex hull of a chain. On the other hand, f lifts uniquely to a monotone map $f_\square: P_\square \rightarrow Q_\square$. Then $|f_\square|$ is 1-Lipschitz (globally), and in particular, uniformly continuous. Since $q: |Q_\square| \rightarrow |Q|$ is uniformly

continuous, so is the composite arrow in the commutative diagram

$$\begin{array}{ccc} |P_{\sqcup}| & \xrightarrow{f_{\sqcup}} & |Q_{\sqcup}| \\ p \downarrow & & q \downarrow \\ |P| & \xrightarrow{f} & |Q|. \end{array}$$

By Theorem 3.1(a), $p: |P_{\sqcup}| \rightarrow |P|$ is a quotient map; in other words, the uniformity of $|P|$ is final with respect to p . Hence $|f|$ is uniformly continuous. We call it the *geometric realization* of f .

It is easy to see that geometric realization of posets and of monotone maps determines a functor (also called the *geometric realization*) from the category of posets and monotone maps to the category of metrizable uniform spaces and uniformly continuous maps.

Theorem 3.8. *The geometric realization functor preserves pullbacks, as well as those pushouts that remain such upon barycentric subdivision.*

Proof. The assertion is equivalent to the preservation of finite products, finite coproducts (which always remain finite coproducts upon barycentric subdivision), embeddings, and those quotient maps that remain quotient maps upon barycentric subdivision. Finite products were considered in Theorem 2.10(a) and embeddings in Corollary 2.7. The preservation of finite coproducts is obvious.

Finally, let $f: P \rightarrow Q$ be a quotient map of posets such that $f^b: P^b \rightarrow Q^b$ is also a quotient map. In particular⁴, f^b is surjective, so every chain in Q is the image of a chain in P . Then $|f_{\sqcup}|: |P_{\sqcup}| \rightarrow |Q_{\sqcup}|$ is a uniformly continuous retraction, and therefore a quotient map. By Theorem 3.1(a), also $q: |Q_{\sqcup}| \rightarrow |Q|$ is a quotient map. Then the composite arrow in the preceding commutative diagram is a quotient map. The assertion now follows from the fact that if a composition $X \rightarrow Y \xrightarrow{f} Z$ is a quotient map, then so is f . $\square$

Theorem 3.8 yields an alternative proof of Theorem 2.10(b,c):

Corollary 3.9. *If P and Q are posets, then $|P * Q|$ and $|P + Q|$ are uniformly homeomorphic to $|P| * |Q|$.*

Proof. $P * Q$ is the pushout of the diagram $P \times Q \times I \supset P \times Q \times \partial I \rightarrow P \times \{\{0\}\} \sqcup Q \times \{\{1\}\}$, where $I = \Delta^{\{0,1\}}$; $P + Q$ is the pushout of the diagram $P \times Q \times [2] \supset P \times Q \times (\{1\} \sqcup \{2\}) \rightarrow P \times \{1\} \sqcup Q \times \{2\}$, where $[2] = (\{1, 2\}, \leq)$; and $X * Y$ is the pushout of the diagram $X \times Y \times [0, 1] \supset X \times Y \times \{0, 1\} \rightarrow P \times \{0\} \sqcup Q \times \{1\}$. Since $|I|$ and $|[2]|$ are uniformly homeomorphic to $[0, 1]$, the assertion follows from Theorem 3.8. $\square$

4. SECOND CANONICAL NEIGHBORHOOD

Lemma 4.1. *Let K be a poset.*

⁴In fact, if the simplicial map f^b is surjective, then f^b and f are quotient maps.

(a) $|MC(r_K)|$ is uniformly homeomorphic to $|MC(\text{id}_K)|$ by a homeomorphism that is the identity on K and extends the homeomorphism $|h(K)| \rightarrow |K^\#| \rightarrow |K|$ given by 2.8 and 3.2.

(b) $|MC^*(r_K)|$ is uniformly homeomorphic to $|MC(\text{id}_{h(K)})|$ by a homeomorphism that is the identity on $|h(K)|$ and extends the homeomorphism $|K| \rightarrow |K^\#| \rightarrow |h(K)|$ given by 2.8 and 3.2.

Beware that $MC^*(r_K)$ is generally not a poset since r_K is generally not open. Note that $MC^*(r_K) \simeq (MC(r_K^*))^*$, where r_K^* is the composition $K^\# \xrightarrow{j_K} (K^*)^\# \xrightarrow{\#} K^*$.

(a). We define $f: |MC(r_K)| \rightarrow |MC(\text{id}_K)|$ as required on the top and bottom, and extend it linearly to the convex hull of every chain. A chain of $MC(\text{id}_K) = K \times [2]$ is of the form $B + A$, where $A = (\alpha_1 < \dots < \alpha_n)$ is a chain in the domain, and $B = (\beta_1 < \dots < \beta_m)$ is a chain in the range, with $\beta_m \leq \alpha_1$. From the similar description of chains of $MC(\text{id}_{h(K)})$ we deduce that a chain of $MC(r_K)$ is of the form $D + C$, where C is a chain of $h(K)$ of the form $[\sigma_1, \tau_1]^* < \dots < [\sigma_r, \tau_r]^*$ and D is a chain of K of the form $\rho_1 < \dots < \rho_s$, with $\rho_s \leq \sigma_1$. It is easy to see that f sends $|h(B) + A|$ onto $|B + A|$ via the join of the uniform homeomorphism $|h(B)| \rightarrow |B|$ with $\text{id}_{|A|}$. It follows that f is a uniform homeomorphism. $\square$

(b). We define $f: |MC^*(r_K)| \rightarrow |MC(\text{id}_{h(K)})|$ as required on the top and bottom, and extend it linearly to the convex hull of every chain. A chain of $MC^*(\text{id}_{h(K)}) = h(K) \times [2]$ is of the form $A + B$, where $A = (\alpha_1 < \dots < \alpha_n)$ is a chain in the domain, and $B = (\beta_1 < \dots < \beta_m)$ is a chain in the range, with $\alpha_n \leq \beta_1$. It follows that a chain of $MC(r_K)$ is of the form $C + D$, where C is a chain of $h(K)$ of the form $[\sigma_1, \tau_1]^* < \dots < [\sigma_r, \tau_r]^*$ and D is a chain of K of the form $\rho_1 < \dots < \rho_s$, with $[\sigma_r, \tau_r]^* \leq [\rho_1, \rho_s]^*$. It is easy to see that f sends $|C + D|$ onto $|C + h(D)|$ via the join of $\text{id}_{|C|}$ and the uniform homeomorphism $|D| \rightarrow |h(D)|$. It follows that f is a uniform homeomorphism. $\square$

Remark 4.2. Similarly to the proof of (a), $|MC(\text{id}_{h(K)})|$ is uniformly homeomorphic to $|MC(r_K)|$ by a homeomorphism that is the identity on $|h(K)|$ and extends the homeomorphism $|h(K)| \rightarrow |K^\#| \rightarrow |K|$ given by 2.8 and 3.2.

Theorem 4.3. *Let K be a poset and L a closed subposet of K . Then $|L|$ is a uniform neighborhood retract of $|K|$.*

Proof. Clearly $|[h(L)]_{h(K)}|$ is a uniform neighborhood of $|h(L)|$ in $|h(K)|$. By 2.8 and 3.2, it corresponds to a uniform neighborhood of $|L|$ in $|K|$. On the other hand, let $\partial N(L) = [h(L)] \setminus h(L) \subset h(K)$, and let $N(L) = L \cup MC^*(r_K|_{\partial N(L)}) \subset MC^*(r_K)$. The monotone map $MC^*(r_K) \rightarrow MC^*(\text{id}_K) \rightarrow K$ is a retraction, and sends $N(L)$ into L ; hence it restricts to a retraction $N(L) \rightarrow L$. Thus $|L|$ is a uniform retract of $|N(L)|$.

Let f be the composition of the uniform homeomorphism $|MC^*(r_K)| \rightarrow |MC^*(\text{id}_{h(K)})|$ in Lemma 4.1(b), with the geometric realization of the projection $\pi: MC^*(\text{id}_{h(K)}) \rightarrow h(K)$. Then f restricts to the identity on $|\partial N(L)|$ and to the uniform homeomorphism $|L| \rightarrow |h(L)|$ on $|L|$. A chain of $MC^*(r_K|_{\partial N(L)})$ is of the form $C + D$, where C is a chain

of $\partial N(L) \subset h(K)$ of the form $[\sigma_1, \tau_1]^* < \cdots < [\sigma_r, \tau_r]^*$ and D is a chain of $L \subset K$ of the form $\rho_1 < \cdots < \rho_s$, with $[\sigma_r, \tau_r]^* \leq [\rho_1, \rho_s]^*$. Then for every chain E of $h(D)$, $C + E$ is a chain of $h(K)$. Since $\partial N(L)$ is disjoint from $h(L)$, it follows that f sends $|C + D|$ homeomorphically onto $|C + h(D)|$ via the join of $\text{id}_{|C|}$ and the uniform homeomorphism $|D| \rightarrow |h(D)|$. It follows that f restricts to a uniform homeomorphism between $|N(L)|$ and $|h(L) \upharpoonright_{h(K)}|$. Since it sends $|L|$ onto $|h(L)|$, we obtain that $|h(L)|$ is a uniform retract of $|h(L) \upharpoonright_{h(K)}|$. $\square$

Remark 4.4. The proof of Theorem 4.3 breaks down if the poset $MC(r_K)$ is used instead of the preposet $MC^*(r_K)$. Indeed, a chain of $MC(r_K|_{\partial N(L)})$ is of the form $D + C$, where C is a chain of $\partial N(L) \subset h(K)$ of the form $[\sigma_1, \tau_1]^* < \cdots < [\sigma_r, \tau_r]^*$ and D is a chain of $L \subset K$ of the form $\rho_1 < \cdots < \rho_s$, with $\rho_s \leq \sigma_1$. Then for a chain E of $h(D)$, $E + C$ is generally not a chain of $h(K)$.

We could still try to follow the argument and see what happens. So let $N'(L) = L \cup MC(r_K|_{\partial N(L)}) \subset MC(r_K)$ and let f' be the composition of the uniform homeomorphism $|MC(r_K)| \rightarrow |MC(\text{id}_{h(K)})|$ in Remark 4.2 with the geometric realization of the projection $\pi: MC(\text{id}_{h(K)}) \rightarrow h(K)$. Suppose that we have two disjoint chains $D + C$ and $D' + C'$ in $N'(L)$, where C and D are as above, and C' is a chain of $\partial N(L) \subset h(K)$ of the form $[\sigma'_1, \tau'_1]^* < \cdots < [\sigma'_r, \tau'_r]^*$ and D is a chain of $L \subset K$ of the form $\rho'_1 < \cdots < \rho'_s$, where $\rho_s < \rho'_1$, $\rho'_s < \sigma'_1$ and $\tau_1 < \sigma'_1$. Then $f'(|D + C|)$ and $f'(|D' + C'|)$ may intersect; so f' restricted to $|N'(L)|$ is generally not injective.

4.5. Relative canonical subdivisions. Let K be a poset and let L be a closed subposet of K . In the notation of the proof of Theorem 4.3, let $h(K, L)$ denote $N(L) \cup (h(K) \setminus h(L))$. Then by the proof of Theorem 4.3, $|h(K, L)|$ is uniformly homeomorphic to $|h(K)|$ and hence to $|K|$.

Dually, let $K_L^\# = MC^*(r_K^*) \setminus (|L| \cup |(K \setminus L)^\#|)$. This contains $L^\#$ and $K \setminus L$, and $|K_L^\#|$ is uniformly homeomorphic to $|K|$ similarly to the above (using part (a) of Lemma 4.1). Note that $(P + Q)^\# \simeq (P^* * Q)_{P^* * \emptyset \cup \emptyset * Q}^\#$, which yields an alternative proof that $|P + Q|$ is uniformly homeomorphic to $|P * Q|$.

4.6. Homotopy completeness. We recall from [13] that a uniform space is called *homotopy complete* if there exists a homotopy $h_t: \overline{|P|} \rightarrow \overline{|P|}$, where $\overline{|P|}$ is the completion of $|P|$, with $h_0 = \text{id}$ and $h_t(\overline{|P|}) \subset |P|$ for $t > 0$.

Lemma 4.7. *Let P be a countable poset. Then $|P|$ is homotopy complete.*

The proof is an extension of the proof that $q_{00} = |2_w^\mathbb{N}|^\bullet$ is homotopy complete, which was given in [13; Corollary 4.11(b)].

Proof. $|P|$ is uniformly homeomorphic to $|P^\#|$, and $P^\#$ is atomic. Hence we may assume without loss of generality that P is atomic.

Let R be the composition $MC(r_P) \rightarrow MC(\text{id}_P) \rightarrow P$ (extending the map r_P), and let $H: |MC(r_P)| \rightarrow |MC(\text{id}_P)|$ be the uniform homeomorphism of Lemma 4.1(a). Define $h_P: \overline{|P|} \times I \rightarrow \overline{|P|}$ to be the unique extension of $|R|H^{-1}$ over the completion, where

$|[2]|$ is identified with $I = [0, 1]$ by the affine homeomorphism sending $\{2\}$ to 1. Further let $h_P^\bullet$ be defined similarly to h_P but using *atomic* geometric realizations throughout, provided that P is either atomic or the dual cone over an atomic poset. Then it is easy to check that $h_{2_w^\bullet}$ coincides with the homotopy h_t constructed in the proof of [13; Corollary 4.11(b)]. On the other hand, the definition of $|P|$ is based on the embedding of $P = (\mathcal{P}, \leq)$ in $\Delta_w^\mathcal{P}$, and h_P is the restriction of $h_{\Delta_w^\mathcal{P}}^\bullet$, which in turn is the restriction of $h_{2_w^\bullet}^\bullet$. Hence $h_P(\overline{|P|} \times (0, 1]) \subset |2_w^\mathcal{P}|^\bullet \cap \overline{|P|} = |P|$. $\square$

Lemma 4.8. $|2_w^\mathbb{N}|$ is a uniform AR.

Proof. By Theorem 2.6, $|2_w^\mathbb{N}|$ is isometric to $|2_w^\mathbb{N}|^\bullet = q_{00}$, which is known to be a uniform AR [13; Corollary 4.11(b)]. $\square$

Remark 4.9. Since $2_w^\mathbb{N} \simeq C^*(\Delta_w^\mathbb{N})$, by Theorem 2.10(b) (or alternatively by Corollary 3.9), $|2_w^\mathbb{N}|$ is uniformly homeomorphic to a cone, and therefore is uniformly contractible. Thus asserting that it is a uniform AR is equivalent (see [13; Lemma 4.32]) to asserting that it is a uniform ANR.

Theorem 4.10. If P is a simplicial complex, then $|P|$ is a uniform ANR.

The finite-dimensional case is due to Isbell [8; 1.9], [10; VI.15].

It is easy to see that the geometric realizations of simplicial complexes are cubohedra in the sense of [13], and so Theorem 4.10 is a special case of [13; Corollary 4.28]. This results in a proof of Theorem 4.10 based on the uniform version of Hanner's criterion of ANR'ness [13; Theorem 4.23(b)], which is in turn inherently rooted in an infinite process.

We now give an alternative proof of Theorem 4.10, which is arguably more technical, but on the other hand more “combinatorial” in that it certainly does not involve any infinite process.

Proof. By [12; Theorem 2.28], P is isomorphic to a subcomplex of the simplex $\Delta_w^\mathbb{N}$. Hence $P^\#$ is isomorphic to a subcomplex of $(\Delta_w^\mathbb{N})^\#$, which in turn is a subcomplex of $(C^*\Delta_w^\mathbb{N})^\# = (2_w^\mathbb{N})^\#$. By Theorem 4.3, we have that $|P^\#|$ is a uniform neighborhood retract of $|(2_w^\mathbb{N})^\#|$. Hence by Corollary 3.2, $|P|$ is a uniform neighborhood retract of $|2_w^\mathbb{N}|$. Since $|2_w^\mathbb{N}|$ is a uniform ANR by Lemma 4.8, so is $|P|$. $\square$

Example 4.11. Consider the poset $[n] = (\{1, \dots, n\}, \leq)$. Let C_n be the amalgamated union of $C^\#$ for all proper subchains $C \subsetneq [n]$. The canonical map $j_n: C_n \rightarrow [n]^\#$ is an injection, but not an embedding for $n > 1$. Consider the map $j := \sqcup_{n \in \mathbb{N}} j_n$, injecting $P := \sqcup_{n \in \mathbb{N}} C_n$ into $Q := \sqcup_{n \in \mathbb{N}} [n]^\#$.

Then the generalized geometric realization $|P|_j$ is not a uniform ANR. Indeed, it follows from Corollary 3.2 that each $|C_n|_{j_n}$ is uniformly homeomorphic to $X_n := \bigcup_{C \subsetneq [n]} |C| \subset |[n]|$. Now $|[n]| = \{(x_0, \dots, x_n) \mid 0 = x_0 \leq \dots \leq x_n = 1\}$, and X_n consists of all $(x_0, \dots, x_n) \in |[n]|$ such that $x_i = x_{i+1}$ for some i . But each $(x_0, \dots, x_n) \in |[n]|$ satisfies $x_{i+1} - x_i \leq \frac{1}{n}$ for some i (by the pigeonhole principle). Hence the $\frac{1}{2n}$ -neighborhood of

X_n in $|[n]|$ is the entire $|[n]|$. Consequently, for each $\varepsilon > 0$ the ε -neighborhood of $|P|_j$ in $|Q|$ contains $|\bigsqcup_{n \in \mathbb{N} \setminus [m]} [n]^\#|$ for some m , and so does not retract uniformly or even continuously onto $|P|_j$.

The following is a direct consequence of [12; Theorem 2.70]

Theorem 4.12. *Let $f: P \rightarrow Q$ be a monotone map between conditionally complete posets. Then $TMC(f)$ is a conditionally complete poset, and $|TMC(f)|$ uniformly strongly deformation retracts onto $|LMC(f)|$.*

Proposition 4.13. *Let $f: P \rightarrow Q$ be a monotone embedding between posets. Then $|\langle MC(f) \rangle|$ uniformly deformation retracts onto $|MC(f)|$.*

Proof. Since f is an embedding, we may identify $\langle MC(f) \rangle$ with a subposet of $Q \times [2]$. Then $h(\langle MC(f) \rangle)$ gets identified with a subposet of $h(Q \times [2]) \simeq h(Q \times [2]^*)$. It is easy to see that the monotone map $r_{Q \times [2]^*}: h(Q \times [2]^*) \rightarrow Q \times [2]^*$ (see [12; 2.79]) sends $h(\langle MC(f) \rangle)$ onto $MC^*(f)$. Using this, similarly to the proof of Theorem 4.3 one constructs a uniform retraction of $|h(\langle MC(f) \rangle)|$ onto $|h(MC^*(f))|$, and by using Lemma 4.1 or similarly to the proof of Lemma 4.7 one constructs a uniform homotopy from this retraction to the identity. $\square$

4.14. Huge mapping cylinder. Let $f: P \rightarrow Q$ be a monotone map between countable posets. Let $j_P: P \hookrightarrow 2^\mathbb{N}$ be the usual embedding $p \mapsto [p]$, where the underlying set of P is identified with a subset of $\mathbb{N}$. Let F be the composition $P \times 2^\mathbb{N} \rightarrow P \xrightarrow{f \times j_P} Q \times 2^\mathbb{N}$ of the projection and the joint map. Finally let $HMC(f)$ be the transitive closure $\langle MC(F) \rangle$. Note that $HMC(f)$ contains canonical copies of $P = P \times \{\emptyset\}$ and $Q = Q \times \{\emptyset\}$.

Corollary 4.15. *Let $f: P \rightarrow Q$ be a monotone map between countable posets. Then $|HMC(f)|$ is uniformly homotopy equivalent to $|MC(f)|$ relative to $|P \sqcup Q|$. If additionally P and Q are conditionally complete posets and f preserves infima, then $HMC(f)$ is a conditionally complete poset.*

5. UNIFORM LOCAL CONTRACTIBILITY

Theorem 5.1. *If P is a countable poset, then $|P|$ is uniformly locally contractible.*

Proof. Given $x, y \in |P|$ with $d(x, y) < 1$, the proof of Theorem 3.1 above produces $x', y' \in |P|$ such that each of the pairs $\{x, x'\}$, $\{x', y'\}$, $\{y', y\}$ lies in the convex hull of a single chain of P , and $d(x, x')$ and $d(y, y')$ are bounded above by $2d(x, y)$. We shall modify this pair of discontinuous maps $(x, y) \mapsto x'$, $(x, y) \mapsto y'$ into a pair of uniformly continuous maps φ, ψ from the uniform neighborhood $\{(x, y) \mid d(x, y) < \delta\}$ of the diagonal in $|P| \times |P|$ into $|P|$ such that $d(x, \varphi(x, y))$ and $d(y, \psi(x, y))$ are bounded above by $\frac{\varepsilon}{2}$. Given δ -close maps $f, g: X \rightarrow |P|$, we then define a homotopy $h_t: X \rightarrow |P|$ by $h_0 = f$, $h_1 = g$, $h_{1/3}(x) = \varphi(f(x), g(x))$ and $h_{2/3}(x) = \psi(f(x), g(x))$ and by linear extension to the remaining values of t . Then $h_{1/3}$ and $h_{2/3}$ are uniformly continuous as compositions of uniformly continuous maps, and are $\frac{\varepsilon}{2}$ -close to h_0 and h_1 , respectively.

Since each of h_1 , $h_{2/3}$ and $h_{1/3}$ is $(\frac{\varepsilon}{2} + \delta)$ -close to h_0 , we infer that h_t is a uniformly continuous $(\frac{\varepsilon}{2} + \delta)$ -homotopy.

It remains to construct φ and ψ . Pick some $x, y \in |P|$ with $d(x, y) < \delta$, and let $A \subset P$ and $B \subset P$ be some finite chains whose convex hulls contain x and y respectively. Arguing as in the proof of Theorem 3.1, we may enlarge P to $\hat{P} = C^*CP$ and consider the unique isomorphisms $a: [m] \rightarrow \hat{A} \subset \hat{P}$ and $b: [n] \rightarrow \hat{B} \subset \hat{P}$. Thus $a(1) = \hat{0} = b(1)$ and $a(m) = \hat{1} = b(n)$. Let $(k_1, l_1), \dots, (k_r, l_r)$ and $(k'_1, l'_1), \dots, (k'_r, l'_r)$, and Z and Z' , and α_i, β_i be as in the proof of Theorem 2.6. We recall that $\alpha_i = x(s)$ for any $s \in a(i) \setminus a(i-1)$, $1 < i \leq m$, and $\beta_i = y(s)$ for any $s \in b(i) \setminus b(i-1)$, $1 < i \leq n$.

The basic problem with the original construction of x', y' in the proof of Theorem 3.1 is that they depend on the choice of A, B . But they should not if φ and ψ are to be continuous; indeed, if A, B are taken to be the *smallest* chains whose convex hulls contain x, y respectively, then a pair $(\tilde{x}, \tilde{y})$ arbitrarily close to (x, y) can give rise to a different pair of chains $(\tilde{A}, \tilde{B})$.

Let δ be such that $\delta \leq \frac{\varepsilon}{6}$ and $N := \frac{1}{4\delta} \in \mathbb{Z}$. Let $u_i, u'_i \in [m]$ and $v_i, v'_i \in [n]$ be the maximal numbers such that $\alpha_{u_i} \geq 1 - 4i\delta$, $\alpha_{u'_i} \geq 1 - (4i+1)\delta$, $\beta_{v_i} \geq 1 - (4i+2)\delta$ and $\beta_{v'_i} \geq 1 - (4i+3)\delta$. Thus $\alpha_{u_0} = 1$ and $u_0 \geq 2$, whereas $\alpha_{u_N} = \alpha_{u_{N-1}} = 0$ and $u_N = m$. It is easy to see⁵ that for each $\kappa \in [m]$ there exists a $\lambda \in [n]$ such that $a(\kappa) \leq b(\lambda)$ and $\beta_\lambda \geq \alpha_\kappa - \delta$. Similarly, for each $\lambda \in [n]$ there exists a $\kappa \in [m]$ such that $b(\lambda) \leq a(\kappa)$ and $\alpha_\kappa \geq \beta_\lambda - \delta$. Hence each $a(u'_i) \leq b(v_i)$ and each $b(v'_i) \leq a(u_{i+1})$. Thus we get an ‘intermediary’ chain C consisting of:

$$\begin{aligned} \hat{0} &\leq a(u_1) \leq a(u_1 + 1) \leq \dots \leq a(u'_1) \\ &\leq b(v_1) \leq b(v_1 + 1) \leq \dots \leq b(v'_1) \\ &\leq a(u_2) \leq a(u_2 + 1) \leq \dots \leq a(u'_2) \\ &\leq \dots \end{aligned}$$

(It should be noted that if $u_i \in Z$, then the inequalities $u_i \leq u'_i \leq \dots \leq u_{i+k} \leq u'_{i+k}$ may all happen to be equalities for an arbitrarily large k . This is the only way that it can happen, for it is easy to see⁶ that if $a(\kappa) \notin B$, then $\alpha_{\kappa+1} \geq \alpha_\kappa - 2\delta$.)

If we use this chain C to construct x' and y' as in the proof of Theorem 3.1, the result will no longer depend on the choice of A and B . However, the definition of C now involves the maximum function, which is discontinuous; so an arbitrarily small change in

⁵Indeed, let i be the minimal number satisfying $k_i \geq \kappa$, and let $\lambda = l_i$. If $i = 1$ then $\beta_\lambda = \alpha_\kappa = 1$. Else $\kappa > k_{i-1}$, hence $\beta_\lambda = \beta_{l_i} \geq \alpha_{k_{i-1}+1} - d(x, y) \geq \alpha_\kappa - \delta$. (The first inequality was established in the proof of Theorem 2.6.)

⁶Let i be the minimal number satisfying $k_i \geq \kappa$. By the hypothesis $i \neq 1$. Then $\kappa > k_{i-1}$, hence $\alpha_\kappa \leq \alpha_{k_{i-1}+1} \leq \beta_{l_i} + d(x, y)$. Next let j be the minimal number satisfying $l'_j \geq l_i$. Then $j \neq 1$ due to $i \neq 1$. Hence $l_i > l'_{j-1}$, so $\beta_{l_i} \leq \beta_{l'_{j-1}+1} \leq \alpha_{k'_j} + d(x, y)$. Thus $\alpha_\kappa \leq \alpha_{k'_j} + 2\delta$. Now if $k'_j > k_i$, then $k'_j > \kappa$ due to $k_i \geq \kappa$. If $k'_j \leq k_i$, then $a(k_i) \leq b(l_i) \leq b(l'_j) \leq a(k'_j) \leq a(k_i)$, implying $k'_j = k_i$ and $i \in Z$. The latter implies $k_i \neq \kappa$ in view of the hypothesis. Then $k_i > \kappa$, and so $k'_j > \kappa$ once again. Thus we obtain that $\alpha_{k'_j} \leq \alpha_{\kappa+1}$.

the coordinates of x can lead to a significant (even though bounded above by δ) change in the coordinates of x' .

Thus we need a new construction of x' and y' that would compensate for the discontinuity of the maximum function. We set $x'(s) = 1 - 4i\delta$ for all $s \in a(u_i) \setminus a(u'_{i-1})$, and (not entirely symmetrically) $y'(s) = 1 - 4i\delta$ for all $s \in b(v_i) \setminus b(v'_{i-1})$. We shall define $x'(s)$ and $y'(s)$ for the remaining values of s by distributing the total jump value 4δ (between e.g. $x'(s)$ and $x'(t)$ for $s \in a(u_i) \setminus a(u'_{i-1})$ and $t \in a(u_{i+1}) \setminus a(u'_i)$, provided that such s and t exist) over all the jumps so as to best approximate the (continuous) uniform distribution. Thus the jump value over $a(j) \setminus a(j-1)$ must be proportional to the step length $\alpha_{j-1} - \alpha_j$ for each $j \in [u_i + 1, u'_i + 1]$. The total horizontal length of the stairs is δ (from $1 - 4i\delta$ to $1 - (4i+1)\delta$, for instance). Therefore we set $x'(s) = (1 - 4i\delta) - 4((1 - 4i\delta) - \alpha_j)$ for all $s \in a(j) \setminus a(j-1)$, for each $j \in [u_i + 1, u'_i]$. Similarly (but not entirely symmetrically) $y'(s) = (1 - 4i\delta) - 4((1 - (4i+2)\delta) - \beta_j)$ for all $s \in b(j) \setminus b(j-1)$, for each $j \in [v_i + 1, v'_i]$. We define $\varphi(x, y) = x'$ and $\psi(x, y) = y'$. We also define $\alpha'_j = x'(s)$ for all $s \in a(j) \setminus a(j-1)$, where $2 \leq j \leq m$, and $\beta'_j = y'(s)$ for all $s \in b(j) \setminus b(j-1)$, where $2 \leq j \leq n$ (beware that this notation is not entirely analogous to that in the proof of Theorem 3.1). Then $\alpha'_2 = \alpha'_{u_0} = 1$ and $\alpha'_{m-1} = \alpha'_{u_N} = 0$, where $u_0 \geq 2$ and $u_N = m$, so $x' \in |A|_a$. Due to the non-symmetric definition of y' , also $\beta'_2 = \beta'_{v_0} = 1$ and $\beta'_{n-1} = \beta'_{v_{N-1}} = 0$, where $v_0 \geq 2$ and $v_{N-1} \leq n$, so $y' \in |B|_b$.

It is easy to check that x' and y' do not depend on the choice of A and B . When $s \in a(u_i) \setminus a(u'_{i-1})$ we have $x(s) \in [\alpha_{u_i}, \alpha_{u'_{i-1}+1}] \subset [1 - 4i\delta, 1 - (4i-3)\delta]$ whereas $x'(s) = 1 - 4i\delta$. When $s \in a(u'_i) \setminus a(u_i)$ we have $x'(s) - D = 4[x(s) - D]$, where $D = 1 - 4i\delta$, so $x'(s) - x(s) = [x'(s) - D] - [x(s) - D] = 3[x(s) - D] \in [0, 3\delta]$. In both cases $|x'(s) - x(s)| \leq 3\delta \leq \varepsilon/2$ as desired. Similarly (but not entirely symmetrically) $|y'(s) - y(s)| \leq 3\delta \leq \varepsilon/2$.

It remains to verify that φ and ψ are uniformly continuous, that is, for each $\zeta > 0$ there exists an $\eta > 0$ such that $d(x, \tilde{x}) < \eta$ and $d(y, \tilde{y}) < \eta$ imply $d(x', \tilde{x}') < \zeta$ and $d(y', \tilde{y}') < \zeta$. By the proof of Theorem 3.1 for each $\theta > 0$ there exists an $\eta > 0$ (namely, $\eta = \theta/5$) such that given $x, \tilde{x} \in |P|$ with $d(x, \tilde{x}) \leq \eta$, there exist $x^*, \tilde{x}^* \in |P|$ such that each of the pairs $\{x, x^*\}$, $\{x^*, \tilde{x}^*\}$, $\{\tilde{x}^*, \tilde{x}\}$ has diameter at most θ and lies in the convex hull of some chain of P . Given $y, \tilde{y} \in |P|$ with $d(y, \tilde{y}) \leq \eta$, we similarly get $y^*, \tilde{y}^*$. Therefore it suffices to consider the case where the pairs $\{x, \tilde{x}\}$ and $\{y, \tilde{y}\}$ lie in the convex hulls of some chains A and B , respectively. Since φ and ψ are well-defined, we may assume that $x', \tilde{x}', y', \tilde{y}'$ are all defined using these A and B . In this case, we set $\eta = \min(\zeta/4, \delta/2)$.

Thus suppose that $d(x, \tilde{x}) < \eta$. In other words, $|\alpha_j(s) - \tilde{\alpha}_j(s)| < \eta$ for all $s \in \mathcal{P}$. Fix some j ; by symmetry we may assume that $\alpha_j(s) \geq \tilde{\alpha}_j(s)$. Since $\eta < \delta$, one of the following four cases has to occur for some i :

- (i) $1 - 4i\delta > \alpha_j(s) \geq \tilde{\alpha}_j(s) \geq 1 - (4i+1)\delta$;
- (ii) $1 - (4i-3)\delta > \alpha_j(s) \geq \tilde{\alpha}_j(s) \geq 1 - 4i\delta$;
- (iii) $1 - (4i-3)\delta > \alpha_j(s) \geq 1 - 4i\delta > \tilde{\alpha}_j(s) \geq 1 - (4i+1)\delta$;

$$(iv) \ 1 - 4i\delta > \alpha_j(s) \geq 1 - (4i + 1)\delta > \tilde{\alpha}_j(s) \geq 1 - (4i + 4)\delta.$$

In the case (i), we have $\alpha'_j(s) - \tilde{\alpha}'_j(s) = 4(\alpha_j(s) - \tilde{\alpha}_j(s))$. In the case (ii), both $\alpha'_j(s)$ and $\tilde{\alpha}'_j(s)$ equal $1 - 4i\delta$. In the case (iii), $\alpha'_j(s) = 1 - 4i\delta$, whereas $\tilde{\alpha}'_j(s) \in [1 - 4i\delta, 1 - 4i\delta - 4\eta]$. Similarly, in the case (iv), $\tilde{\alpha}'_j(s) = 1 - (4i + 4)\delta$, whereas $\alpha'_j(s) \in [1 - (4i + 4)\delta + 4\eta, 1 - (4i + 4)\delta]$.

In all cases, $\alpha'_j(s) - \tilde{\alpha}'_j(s) \in [0, 4\eta]$. This shows that $d(x', \tilde{x}') \leq 4\eta$. Thus φ is uniformly continuous; the uniform continuity of ψ is verified similarly. $\square$

Example 5.2. Given a preposet $P = (\mathcal{P}, \leq)$, we define the “co-deleted prejoin” $P \boxplus P^*$ to be the preposet $(\mathcal{P} \sqcup \mathcal{P}^*, \preceq)$, where $\mathcal{P}^* = \{p^* \mid p \in P\}$ is a just fancy notation for a copy of $\mathcal{P}$, and the relation is defined by

- $p \preceq q$ iff $p \leq q$;
- $p^* \preceq q^*$ iff $p \geq q$;
- $p^* \preceq q$ never holds;
- $p \preceq q^*$ iff either $p \leq q$ or $p \geq q$

for all $p, q \in \mathcal{P}$. Note that $P \boxplus P^*$ need not be a poset even if P is.

Let us define $j: P \rightarrow (P \boxplus P^*)^\#$ by $j(p) = [p, p^*]$. Obviously, j is a monotone embedding, i.e. $p \leq q$ if and only if $j(p) \preceq j(q)$. We claim that $|j|$ is a homotopy equivalence. Indeed, $|j|$ is homotopic to the composition $|P| \xrightarrow{|i|} |P| \xrightarrow{h} |P^\#|$, where h is the uniform homeomorphism. On the other hand, $i^\flat$ is split by the simplicial map $r: (P \boxplus P^*)^\flat \rightarrow P^\flat$, defined on vertices by $p, p^* \mapsto p$. Given a chain $\sigma = (p_1 < \dots < p_n) \in P^\flat$ we have $r^{-1}(\lceil \sigma \rceil) = \lceil \sigma \boxplus \bar{\sigma} \rceil$, where $\sigma \boxplus \bar{\sigma}$ denotes the chain $(p_1 < \dots < p_n < p_n^* < \dots < p_1^*) \in (P \boxplus P^*)^\flat$. Since r is simplicial, it follows that $|r|: |(P \boxplus P^*)^\flat| \rightarrow |P^\flat|$ has contractible point-inverses, and therefore (or by Quillen’s fiber lemma) is a homotopy equivalence. If k is a homotopy inverse to $|r|$, then $k = k|r i^\flat| \simeq |i^\flat|$, so $|i^\flat|$ is also a homotopy equivalence.

Let K_0 be the preposet of the four sets 0 , $\{0, 1\}$, $\{0, 2\}$ and $\{\{0, 1\}, \{0, 2\}\}$ ordered by $\in$. Thus $|K_0|$ is homeomorphic to S^1 . Let $K_{n+1} = K_n \boxplus K_n^*$. Finally let $K = K_0 \sqcup K_1 \sqcup \dots$.

We claim that $|K|$ is not uniformly locally contractible (and in particular is not a uniform ANR). Indeed, by the above we have an embedding $f_n: K_0 \rightarrow K_n^{\#n}$ such that $|f_n|$ is a homotopy equivalence. In order to use the d_∞ metric, which has been shown to work only for posets, we consider the transitive closure. Let f'_n be the composition $K_0 \xrightarrow{f_n} K_n^{\#n} \subset \langle K_n^{\#n} \rangle \subset \langle K_n \rangle^{\#n}$. Since f'_n is monotone, the image of $|f'_n|$ has diameter 1 in the d_∞ metric on $|\langle K_n \rangle^{\#n}|$, hence by Corollary 3.2, the image of the composition $|K_0| \xrightarrow{|f'_n|} |\langle K_n \rangle^{\#n}| \xrightarrow{h_n} |\langle K_n \rangle|$ has diameter 2^{-n} in the d_∞ metric on $|\langle K_n \rangle|$. Since $\text{id}: (|\langle K_n \rangle|, d_\infty) \rightarrow (|\langle K_n \rangle|, d)$ is 1-Lipschitz, the image of the composition $|K_0| \xrightarrow{|f'_n|} |\langle K_n \rangle^{\#n}| \xrightarrow{h_n} |K_n|$ has diameter at most 2^{-n} with respect to the usual metric on $|K_n|$. However this composition is not null-homotopic since it is a homotopy equivalence.

We note that the preposet K in Example 5.2 satisfies the following property (P): For each $\varepsilon > 0$ there exists an essential map $S^1 \rightarrow |K|$ with image of diameter $< \varepsilon$. On the

other hand, since $|K^\flat|$ is a uniform ANR, $|K|$ is a non-uniform ANR, and in particular satisfies the non-uniform homotopy extension property. It follows that every metrizable uniform space that is uniformly homotopy equivalent to $|K|$ satisfies (P) as well. In particular, we get the following

Theorem 5.3. *There exists a countable preposet whose geometric realization is not uniformly homotopy equivalent to a uniform ANR, nor even to a uniformly locally contractible metrizable uniform space.*

6. COMBINATORICS OF COVERS

In this subsection we shall need basic operations and relations on covers as introduced in [13; 2.6], as well as the following additional notation.

6.1. Nerve. We recall that the *nerve* of a cover $C \subset 2^S$ of a set S is the simplicial poset $N(C) \subset 2^C$, where a subset $B \subset C$ is a simplex of $N(C)$ iff $\bigcap B$ (the intersection of all elements of B) is nonempty. The notion of nerve was introduced by Alexandroff [1]. We note that for a cover C ,

- C countable and point-finite iff $N(C)$ is a simplicial complex;
- C is countable and Noetherian iff $N(C)$ is a Noetherian simplicial complex;
- C is countable and star-finite iff $N(C)$ is a locally finite simplicial complex.

6.2. Simplex determined by subset. Given a nonempty $T \subset S$ that is contained in at least one element of C , let $\Delta_C(T)$ denote the element $\{U \in C \mid T \subset U\}$ of $N(C)$. Given an $s \in S$, we write $\Delta_C(s) = \Delta_C(\{s\})$. Note that every element of $N(C)$ belongs to some simplex of $N(C)$ of the form $\lceil \Delta_C(s) \rceil$ for some $s \in S$.

6.3. Cover by open stars. If P is a poset, by the *open star* $\overset{\circ}{\text{st}}(p, P)$ of a $p \in P$ we mean $|\text{st}(p, P)| \setminus |\text{st}(p, P) \setminus \lfloor p \rfloor|$. (We recall that $\text{st}(p, P) = \lceil \lfloor p \rfloor \rceil$.)

If P is atomic with atom set Λ , we have the cover $\{\overset{\circ}{\text{st}}(\lambda, P) \mid \lambda \in \Lambda\}$ of $|P|^\bullet$ by *open stars of vertices*. It is easy to see that the open star of a vertex $v \in \Lambda$ in $|P|^\bullet$ is precisely the set of points of $|P|^\bullet \subset [0, 1]^\Lambda$ whose v th coordinate is nonzero. On the other hand, the set of points of $|P|^\bullet$ whose v th coordinate equals 1 is precisely the dual cone $|\lfloor v \rfloor|$ of v . Thus we get a cover of $|P|^\bullet$ by *dual cones of vertices*, and it follows that the cover of $|P|^\bullet$ by the open stars of vertices is uniform with Lebesgue number $1 - \varepsilon$ for each $\varepsilon > 0$.

6.4. Strict shrinking. If C is a cover of a uniform space X , a *strict shrinking* of C is a cover C' of X such that there exist a uniform cover D of X and a bijection $C \rightarrow C'$, denoted $U \mapsto U'$, such that $\text{st}(U', D) \subset U$ for each $U \in C$.

The following is a strengthened statement of [10; IV.19]:

Lemma 6.5. *Let X be a uniform space, C a uniform cover of X , and D a uniform star-refinement of C . Then $C_D := \{U_D \mid U \in C\}$, where $U_D = X \setminus \text{st}(X \setminus U, D)$, is a strict shrinking of C .*

Beware that C_D is not intended to be a uniform cover here.

Proof. Obviously, $\text{st}(U_D, D) \subset U$ for each $U \in C$. On the other hand, since D is a star-refinement of C , for every $x \in X$, $\text{st}(x, D)$ lies in some $U \in C$, whence $x \in U_D$. Thus C_D is a cover of X . $\square$

Example 6.6. Let P be a simplicial complex or more generally an atomic poset. Let C be the cover of $|P|$ by open stars of vertices, and let $C^\#$ denote the image of the cover of $|P^\#|$ by open stars of vertices under the homeomorphism $|P^\#| \cong |P|$. Then $C^\#$ star-refines C , and $C_{C^\#}$ is the cover of $|P|$ by dual cones of vertices.

Lemma 6.7. *Let X be a metrizable uniform space and C a countable point-finite uniform open cover of X .*

(a) *There exists a uniformly continuous map $\varphi_C: X \rightarrow |N(C)|^\bullet$ that sends each $x \in X$ into the interior of $|\lceil \Delta_C(x) \rceil|^\bullet$.*

(b) *If D is a uniform star-refinement of C , then there exists a uniformly continuous map $\varphi_{C,D}: X \rightarrow |N(C)|^\bullet$ that sends each $x \in X$ into the intersection of $|\lceil \Delta_{C_D}(x) \rceil|^\bullet$ and the interior of $|\lceil \Delta_C(x) \rceil|^\bullet$.*

Note that the conclusion of (a) implies that each $U \in C$ is the preimage of the open star of the vertex $\{U\}$ of $N(C)$. The conclusion of (c) implies additionally that each $U_D \in C_D$ lies in the preimage of the dual cone of the vertex $\{U\}$ of $N(C)$.

Proof. Note that (b) implies (a).

To prove (b), let λ be a Lebesgue number of D . Then each $d(U_D, X \setminus U) \geq \lambda$. For each $U \in C$ define $f_U: X \rightarrow [0, 1]$ by $f_U(x) = \min(\lambda^{-1}d(x, X \setminus U), 1)$. Let us consider $\varphi = \prod f_U: X \rightarrow l_\infty^c$. Then $\{U \in C \mid f_U(x) > 0\} = \Delta_C(x)$, and since C' is a cover, $\{U \in C \mid f_U(x) = 1\} \neq \emptyset$. Hence $\varphi(X) \subset |N(C)|^\bullet$. Next, $f_U^{-1}(0) = X \setminus U$, which implies the assertion on C , and $f_U(U_D) = \{1\}$, which implies the assertion on C_D . Finally, $|f_U(x) - f_U(y)| \leq \lambda^{-1}d(x, y)$ for each $U \in C$, so φ is uniformly continuous. $\square$

6.8. Subordinated map. Let C be a cover of a set S , and $f: S \rightarrow |P|^\bullet$ a map, where P is an atomic poset. We say that f is *subordinated* to C if C is refined by $f^{-1}(D)$, where D is the cover of $|P|^\bullet$ by the open stars of vertices. A homotopy $h_t: S \rightarrow |P|^\bullet$ is said to be subordinated to C if it is through maps subordinated to C .

Lemma 6.9. *Let X be a metrizable uniform space.*

(a) *If C a point-finite countable uniform cover of X , and D, E are uniform star-refinements of C , then $\varphi_{C,D}$ and $\varphi_{C,E}$ are uniformly homotopic by a homotopy subordinated to C .*

(b) *Let P a simplicial complex and $f: X \rightarrow |P|^\bullet$ a uniformly continuous map. Let E be the cover of $|P|^\bullet$ by open stars of vertices, and let $C = f^{-1}(E)$ and $D = f^{-1}(C^\#)$. Then $\varphi_{C,D}: X \rightarrow |N(C)|^\bullet \subset |N(E)|^\bullet$ is uniformly homotopic to $f: X \rightarrow |P|^\bullet = |N(E)|^\bullet$ by a homotopy subordinated to C .*

(c) *Let C be a point-finite countable uniform cover of X and D a uniform star-refinement of C . Let E_D be a subset of C_D that still covers X , and E the corresponding*

subset of C . Then $\varphi_{C,D}: X \rightarrow |N(C)|^\bullet$ is uniformly homotopic to $\varphi_{E,D}: X \rightarrow |N(E)|^\bullet \subset |N(C)|^\bullet$ by a homotopy subordinated to C .

Proof. (a). By considering $\{U_D \cup U_E \mid U \in C\}$ we may assume that each element of C_D is contained in the corresponding element of C_E . Then $\Delta_{C_D}(x) \subset \Delta_{C_E}(x)$ for each $x \in X$. Hence $[\Delta_{C_E}(x), \Delta_C(x)] \subset [\Delta_{C_D}(x), \Delta_C(x)]$. Therefore $\varphi_{C,D}(x)$ and $\varphi_{C,E}(x)$ both belong to $[[\Delta_{C_D}(x), \Delta_C(x)]]^\bullet$. Thus the linear homotopy between $\varphi_{C,D}$ and $\varphi_{C,E}$ in q_{00} has values in $|N(C)|^\bullet$. Since both $\varphi_{C,D}$ and $\varphi_{C,E}$ are subordinated to C , so is the homotopy. $\square$

(b). By Lemma 6.7, $\varphi_{C,D}(x) \in [[\Delta_{C_D}(x), \Delta_C(x)]]^\bullet$ for each $x \in X$. Clearly, $f(x) \in [[\Delta_{C_D}(x), \Delta_C(x)]]^\bullet$ for each $x \in X$. Thus the linear homotopy between f and $\varphi_{C,D}$ in q_{00} has values in $|N(C)|^\bullet$. Since both f and $\varphi_{C,D}$ are subordinated to C , so is the homotopy. $\square$

(c). Given an $x \in X$, we have $\Delta_{E_D}(x) \subset \Delta_{C_D}(x)$ and $\Delta_E(x) \subset \Delta_C(x)$. Hence $\Delta_{C_D}(x), \Delta_E(x) \in [\Delta_{E_D}(x), \Delta_C(x)]$. Therefore $\varphi_{C,D}(x)$ and $\varphi_{E,D}(x)$ both belong to $[[\Delta_{E_D}(x), \Delta_C(x)]]^\bullet$. Thus the linear homotopy between $\varphi_{C,D}$ and $\varphi_{E,D}$ in q_{00} has values in $|N(C)|^\bullet$. Since both $\varphi_{C,D}$ and $\varphi_{E,D}$ are subordinated to C , so is the homotopy. $\square$

6.10. Intersection poset and Venn diagram. Given a cover C of a set S , the *intersection poset* $IP(C)$ is the subposet of 2^C consisting of all nonempty $B \subset C$ such that $\bigcap B$ is not contained in any element of $C \setminus B$. The terminology “intersection poset” derives from Lemma 6.11(a) below, which however characterizes $IP(C)$ only up to isomorphism, and not as a subposet of Δ^C .

The *Venn diagram* $VD(C)$ is the subposet of 2^C consisting of all $B \subset C$ such that $\bigcap B$ is not contained in $\bigcup(C \setminus B)$. This is a formalization of the intuitive notion of a “Venn diagram”, also known as “Euler diagram”, from courses of “abstract mathematics”, for it can be argued that $VD(C)$ contains all the combinatorial information on containment of points of X in elements of C (see Lemma 6.11(b) below) — and nothing else (see Lemma 6.13(b) below).

Clearly, $VD(C) \subset IP(C) \subset N(C)$ and $\lceil VD(C) \rceil = \lceil IP(C) \rceil = N(C)$. Note that if C is countable and point-finite, then $IP(C)$ and $VD(C)$ are cone complexes. Clearly $\Delta_C(x)$ belongs to $VD(C)$ for every $x \in S$; in contrast, $\Delta_C(T)$ belongs to $IP(C)$ for every $T \subset S$ that is contained in at least one element of C .

Lemma 6.11. *Let C be a cover of a set S .*

(a) *$IP(C)$ is isomorphic to the poset consisting of arbitrary nonempty intersections of elements of C , ordered by reverse inclusion. In particular, $IP(C)$ is conditionally complete.*

(b) *$VD(C)$ is isomorphic to the poset consisting of those intersections of elements of C that are of the form $\bigcap \Delta_C(s)$ for some $s \in S$, ordered by reverse inclusion.*

Proof. It will be convenient to work in a slightly greater generality. The definitions of $N(C)$, $IP(C)$ and $VD(C)$ generalize straightforwardly for any collection $\varphi: C \rightarrow 2^S$

of subsets of S (possibly with repeated subsets $\varphi(U) = \varphi(U')$ and with $\bigcup \varphi(C)$ not necessarily covering the whole of S). It is easy to see that

- $N(\varphi) = \Phi^{-1}(\Delta^S)$, where $\Phi: \Delta^C \rightarrow 2^S$ is defined by $D \mapsto \bigcap \varphi(D)$;
- $IP(\varphi) = \Delta_\varphi(\Delta^S) \setminus \{\emptyset\}$, where $\Delta_\varphi: \Delta^S \rightarrow 2^C$ is defined by $T \mapsto \varphi^{-1}(\lfloor T \rfloor)$;
- $VD(\varphi) = \Delta_\varphi(A(\Delta^S)) \setminus \{\emptyset\}$, where the subset $A(\Delta^S) = \{\{s\} \mid s \in S\}$ of Δ^S should not be confused with the element S of Δ^S .

We note that the maps Φ and Δ_φ are anti-monotone, and restrict to mutually inverse bijections between $IP(\varphi)$ and $\Phi(\Delta^C) \setminus \{\emptyset\}$. In particular, $IP(\varphi)$ is isomorphic to $(\Phi(\Delta^C) \setminus \{\emptyset\})^*$, which implies the first assertion of (a). Similarly, $VD(\varphi)$ is isomorphic to $\Phi(\Delta_\varphi(A(\Delta^S)) \setminus \{\emptyset\})^*$, which yields (b). $\square$

Remark 6.12. It follows from the proof that $IP(\varphi) \simeq IP(\Phi)$ and $VD(\varphi) \simeq VD(\Phi)$.

Lemma 6.13. *Let P be a poset embedded in some Δ^Λ . Let C be the cover $\{\lfloor \lambda \rfloor \cap P \mid \lambda \in \Lambda\}$ of the underlying set of P by the dual cones of vertices.*

(a) $VD(C) \simeq P$.

(b) $IP(C) = VD(C)$ if and only if for every $R \subset A(\Delta^\Lambda)$, the set of all upper bounds of R in P either is empty or is the dual cone in P of a single element of P .

Proof. (a). By Lemma 6.11(b), $VD(C)$ is isomorphic to the poset consisting of the dual cones $\lfloor p \rfloor^{\Delta^\Lambda} \cap P = \lfloor p \rfloor^P$ of all elements $p \in P$, ordered by reverse inclusion. The latter is obviously isomorphic to P . $\square$

(b). By Lemma 6.11(a), $IP(C)$ is isomorphic to the poset of all nonempty intersections of the form $\bigcap \sigma$, where $\sigma \subset C$, ordered by reverse inclusion. We have $\bigcap \sigma = P \cap \bigcap_{\lambda \in R_\sigma} \lfloor \lambda \rfloor$, where $R_\sigma = \{\{\lambda\} \mid \lambda \in \Lambda, (\lfloor \lambda \rfloor \cap P) \in \sigma\}$. (The subset R_σ of Δ^Λ should not be confused with the element $\bigcup R_\sigma$ of Δ^Λ .) Thus $IP(C)$ is in bijection with the set of all nonempty intersections of the form $P \cap \lfloor R \rfloor$, where $R \subset \Lambda$. By the proof of Lemma 6.11, the same bijection sends $VD(C)$ onto the set of the dual cones $\lfloor p \rfloor^P$ of all elements $p \in P$, and the assertion follows. $\square$

Corollary 6.14. *Let P be an atomic poset, and let C be the cover of P by the dual cones of its atoms. Then $IP(C) = VD(C)$ if and only if P is conditionally complete.*

It is not hard to see that the same assertion is true of the cover of $|P|$ by the geometric realizations of the dual cones of the vertices of P , and of the cover of $|P|$ by the open stars of these vertices.

Proof. Let us embed P in $\Delta^{A(P)}$ as in [12; Lemma 2.25]. Lemma 6.13(b) then says that $IP(C) = VD(C)$ if and only if every $R \subset A(P)$ that has an upper bound in P has a least upper bound in P . This proves the “if” assertion, and the “only if” assertion now follows from [12; Lemma 2.26]. Alternatively, the “only if” assertion follows from Lemma 6.13(a) and the second assertion of Lemma 6.11(a). $\square$

6.15. Canonical bonding map. Let C and D be covers of a set S , and suppose that C star-refines D . We define a map $\varphi_D^C: N(C) \rightarrow N(D)^\#$ by sending each $\sigma \in N(C)$ into $[\Delta_D(\bigcup \sigma), \Delta_D(\bigcap \sigma)] \in N(D)^\#$. Here $\Delta_D(\bigcup \sigma)$ is non-empty by the star-refinement hypothesis; every vertex of $\Delta_D(\bigcup \sigma)$ is obviously a vertex of $\Delta_D(\bigcap \sigma)$; and $\Delta_D(\bigcap \sigma) \in N(D)$ since $\bigcap \sigma \neq \emptyset$.

Given a $\tau \leq \sigma$, we have $\bigcup \tau \subset \bigcup \sigma$, whence $\Delta_D(\bigcup \tau) \geq \Delta_D(\bigcup \sigma)$; and $\bigcap \tau \supset \bigcap \sigma$, whence $\Delta_D(\bigcap \tau) \leq \Delta_D(\bigcap \sigma)$. Thus φ_D^C is monotone.

Finally, recall that $IP(D)$ contains every element of $N(D)$ of the form $\Delta_D(T)$, where $T \subset S$. Hence $[\Delta_D(\bigcup \sigma), \Delta_D(\bigcap \sigma)]$ belongs to the isomorphic copy of $IP(D)^\#$ in $N(D)^\#$. Thus we may write $\varphi_D^C: N(C) \rightarrow IP(D)^\#$.

Remark 6.16. Let us discuss some motivation/geometry behind the definition of φ_D^C .

If $V \in \sigma$ then $\{V\} \leq \sigma$, so by the above $\Delta_D(V)$ belongs to $[\Delta_D(\bigcup \sigma), \Delta_D(\bigcap \sigma)]$. Hence the map defined on vertices by $\{V\} \mapsto [\Delta_D(V), \Delta_D(V)]$ extends uniquely to a monotone map $N(C) \rightarrow N(D)^\#$ sending each simplex $[\sigma]$ into the cube $[\Delta_D(\bigcup \sigma), \Delta_D(\bigcap \sigma)]$.

If C strongly star-refines D , then $\Delta_D(\text{st}(V, C)) \in N(C)$, and it is not hard to see that φ_D^C sends each $\text{st}(\{V\}, N(C))$ into the canonical subdivision of the dual cone of $\Delta_D(\text{st}(V, C))$ in the subcomplex $\bigcup \{[\Delta_D(x)] \mid x \in V\}$ of $N(D)$.

Proposition 6.17. *Let X be a metrizable uniform space, P and Q simplicial complexes, and $f: X \rightarrow |P|$ and $g: X \rightarrow |Q|$ uniformly continuous maps. Let C_P and C_Q be the covers of $|P|$ and $|Q|$ by the open stars of vertices, and let $D := g^{-1}(C_Q)$. If $C := f^{-1}(C_P)$ refines $E := g^{-1}(C_Q^\#)$, then the composition $X \xrightarrow{f} |P| \xrightarrow{|\varphi_D^C|} |Q|^\# \cong |Q|$ is uniformly homotopic to g by a homotopy subordinated to D .*

Note that C star-refines D , since E does; thus φ_D^C is defined.

Proof. Given an $x \in X$, we have $g(x) \in |[\Delta_{D_E}(x), \Delta_D(x)]|$. On the other hand, $f(x) \in |[\Delta_C(x)]|$. Hence $\varphi_D^C(f(x)) \in |[\Delta_D(\bigcup \Delta_C(x)), \Delta_D(\bigcap \Delta_C(x))]|$. Since $x \in \bigcap \Delta_C(x)$, we have $\Delta_D(\bigcap \Delta_C(x)) \subset \Delta_D(x)$. On the other hand, since C refines E , and $\text{st}(U_E, E) \subset U$ for each $U \in D$, we have that $x \in U_E$ implies $\bigcup \Delta_C(x) \subset U$. Hence $\Delta_{D_E}(x) \subset \Delta_D(\bigcup \Delta_C(x))$. Thus both $g(x)$ and $\varphi_D^C(f(x))$ lie in $[\Delta_{D_E}(x), \Delta_D(x)]$. Then the linear homotopy between g and $\varphi_D^C f$ in q_{00} has values in $|N(D)| \subset |Q|$. Since both g and φ_D^C are subordinated to D , so is the homotopy. $\square$

Proposition 6.18. *Let C and D be covers of a set S . If C strongly star-refines D , there exists a simplicial map $N(C) \xrightarrow{g} N(D)$ such that $N(C)^\# \xrightarrow{g^\#} N(D)^\#$ is homotopic to the composition $N(C)^\# \xrightarrow{(\varphi_D^C)^\#} N(D)^\# \xrightarrow{r^\#} N(D)^\#$, where r stands for either $r_{N(D)}$ or $r_{N(D)}^*$, by a monotone homotopy $N(C)^\# \times I \rightarrow N(D)^\#$ sending each block of the form $[\sigma]^\# \times I$ into the star of $[\Delta_D(\bigcup \sigma), \Delta_D(\bigcap \sigma)]$ in $N(D)^\#$.*

Proof. Given a vertex $\{V\}$ of $N(C)$, the hypothesis furnishes a vertex $\{V'\}$ of $N(D)$ such that $\text{st}(V, C) \subset V'$. For each $\sigma \in N(C)$ and each $V \in \sigma$ we have $\bigcup \sigma \subset V'$, and consequently $\{V'\} \in [\Delta_D(\bigcup \sigma)]$. Hence $V \mapsto V'$ extends to a simplicial map

$g: N(C) \rightarrow N(D)$ sending each σ onto some $\sigma' \subset \Delta_D(\bigcup \sigma) \subset \Delta_D(\bigcap \sigma)$. Then $g^\#$ sends each $[\sigma, \tau]$ onto $[\sigma', \tau']$. The composition $N(C)^\# \xrightarrow{(\varphi_D^C)^\#} N(D)^{\#\#} \xrightarrow{r^\#} N(D)^\#$ sends it onto $[\Delta_D(\bigcup \sigma), \Delta_D(\bigcup \tau)]$ when $r = r_{N(D)}$ and onto $[\Delta_D(\bigcap \sigma), \Delta_D(\bigcap \tau)]$ when $r = r_{N(D)}^*$. The required homotopy is defined by sending each $([\sigma, \tau], \{0, 1\})$ onto $[\sigma', \Delta_D(\bigcup \sigma)]$ in the first case and onto $[\sigma', \Delta_D(\bigcap \sigma)]$ in the second case. $\square$

Example 6.19. Here is a simple example of a canonical bonding map that is neither closed nor infima-preserving.

Let $C = \{U_1, U_2, U_3\}$ be the cover of Δ^2 by the open stars of vertices. Let $D = \{U_1, U_2 \cup U_3, U_1 \cup U_2 \cup U_3\}$. Then C star-refines D . The canonical bonding map $\varphi_D^C: N(C) \rightarrow N(D)^\#$ is the composition of the simplicial surjection $\Delta^2 \rightarrow \Delta^1$ (see [12; Example 2.68]), the diagonal embedding $\Delta^1 \rightarrow \Delta^1 \times \Delta^1$ (see [12; Example 2.14]), and a subcomplex inclusion $\Delta^1 \times \Delta^1 \subset (\Delta^2)^\#$.

7. APPROXIMATION OF MAPS

Theorem 7.1. *Let Q be a countable conditionally complete poset. Then $|Q|$ satisfies the Hahn property.*

Proof. Given an $\varepsilon > 0$, let C' be the cover of $|Q^{\#n}|$ by the open stars of vertices, where $2^{-n+1} < \varepsilon$ and $n \geq 1$ so that $Q^{\#n}$ is atomic. Since Q is conditionally complete, so is $Q^{\#n}$, and therefore $IP(C') = VD(C') \simeq Q^{\#n}$. Let $h_n: |Q^{\#n}| \rightarrow |Q|$ be the uniform homeomorphism given by Corollary 3.2, and let $C = h_n(C')$. Let δ be the Lebesgue number of C with respect to the d_∞ metric on $|Q|$.

Given a metric space X and a (γ, δ) -continuous map $f: X \rightarrow |Q|$ for some $\gamma > 0$, let E be the cover of X by $\frac{\gamma}{4}$ -balls. Then E star-refines $D := f^{-1}(C)$. Let Φ denote the composition

$$X \xrightarrow{\varphi_E} |N(E)| \xrightarrow{|\varphi_D^E|} |IP(D)^\#| \subset |IP(C)^\#| \xrightarrow{h} |IP(C)| \cong |Q^{\#n}| \xrightarrow{h_n} |Q|,$$

where φ_E is the uniformly continuous map given by Lemma 6.7 and φ_D^E is the canonical bonding map. Given an $x \in X$, by Lemma 6.7 $\varphi_E(x) \in |\lceil \Delta_E(x) \rceil|$. By the definition of φ_D^E we have $\varphi_D^E(\Delta_E(x)) = [\Delta_D(\bigcup \Delta_E(x)), \Delta_D(\bigcap \Delta_E(x))] \subset \lceil \Delta_D(\bigcap \Delta_E(x)) \rceil \subset \lceil \Delta_D(x) \rceil$. The latter is identified with $\lceil \Delta_C(f(x)) \rceil$, where $\Delta_C(f(x))$ is an element of $IP(C) \simeq Q^{\#n}$, and it follows that $\Phi(x) \in h_n(|\lceil \Delta_C(f(x)) \rceil|)$. Now $|\lceil \Delta_C(f(x)) \rceil|$ has diameter ≤ 2 with respect to the d_∞ metric on $|Q^{\#n}|$, so $h_n(|\lceil \Delta_C(f(x)) \rceil|)$ has diameter $\leq 2^{-n+1}$ with respect to the d_∞ metric on $|Q|$. Since this set contains both $\Phi(x)$ and $f(x)$, we infer that Φ is ε -close to f with respect to the d_∞ metric on $|Q|$. $\square$

We define a *uniform polyhedron* to be the geometric realization of a (countable) conditionally complete poset. Theorem 5.1, Theorem 7.1, and [13; Theorem 4.22] have the following

Corollary 7.2. *Uniform polyhedra are uniform ANRs.*

Lemma 7.3. *For each $\varepsilon > 0$ there exist an n and a $\delta > 0$ such that for each $\gamma > 0$ there exists an M such that for each $m \geq M$ the following holds. Let P be a preposet and Q a conditionally complete poset, and $f: |P| \rightarrow |Q|$ be a (γ, δ) -continuous map. Then there exists a monotone map $g: P^{\#m} \rightarrow Q^{\#n}$ such that the composition $|P| \xrightarrow{h_m^{-1}} |P^{\#m}| \xrightarrow{g} |Q^{\#n}| \xrightarrow{h_n} |Q|$ is ε -close to f .*

Proof. Let $2^{-n+1} < \varepsilon$, $n \geq 1$, let $\delta < 2^{-n-1}$, and let $2^{-M+1} < \gamma/4$, $M \geq 1$.

Let C' be the cover of $|Q^{\#n}|$ by the open stars of vertices (using that $Q^{\#n}$ is atomic due to $n \geq 1$). Since C' has Lebesgue number $\frac{1}{2}$ with respect to the usual metric d on $|Q^{\#n}|$, it also has Lebesgue number $\frac{1}{2}$ with respect to the d_∞ metric, due to $d(x, y) \leq d_\infty(x, y)$. Then $C := h_n(C')$ has Lebesgue number 2^{-n-1} (and therefore also Lebesgue number δ) with respect to the d_∞ metric on $|Q|$.

Let E' be the cover of $|P^{\#m}|$ by the open stars of vertices (using that $P^{\#m}$ is atomic due to $m \geq M \geq 1$). Then E' refines the cover of $|P^{\#m}|$ by balls of radius 2 about every vertex of $P^{\#m}$ with respect to the d_∞ metric on $|P^{\#m}|$. Hence $E := h_m(E')$ refines the cover of $|P|$ by balls of radius 2^{-m+1} (and therefore also that by balls of radius $\gamma/4$) about all points of $|P|$ with respect to the d_∞ metric on $|P|$. We note that the composition $\varphi: |P| \xrightarrow{h_m^{-1}} |P^{\#m}| \cong |VD(E)| \subset |N(E)|$ satisfies $\varphi(x) \in \lceil \Delta_E(x) \rceil$.

The assertion now follows by the proof of Theorem 7.1. $\square$

From the preceding lemma we infer

Theorem 7.4. *For each $\varepsilon > 0$ there exists an n such that the following holds. Let $f: |P| \rightarrow |Q|$ be a uniformly continuous map, where P is a preposet and Q is a conditionally complete poset. Then there exists an M such that for each $m \geq M$ there exists a monotone map $g: P^{\#m} \rightarrow Q^{\#n}$ such that the composition $|P| \xrightarrow{h_m^{-1}} |P^{\#m}| \xrightarrow{g} |Q^{\#n}| \xrightarrow{h_n} |Q|$ is ε -close to f .*

Example 7.5. Let $P_1 = [2]$ and let $P_{i+1} = P_i + [2]$. Finally let $P = \bigsqcup_{n \in \mathbb{N}} P_{2^n+1}$. We claim that $|P|$ does not satisfy the Hahn property (and in particular is not a uniform ANR).

Indeed, let $Q_n = (P_{2^n+1})^{\#n}$, and let C_n be the cover of $|Q_n|$ by the stars of atoms of Q_n . Then $|VD(C_n)| \cong |Q_n| \cong |P_{2^n+1}|$ is a 2^n -sphere; but we shall now show that $|N(C_n)|$ is contractible.

If K is a poset, then $K^{\#n}$ is isomorphic to the poset consisting of non-decreasing sequences $a = (a_1 \leq \dots \leq a_{2^n})$ of elements of K , where $a \geq b$ iff $a_i \leq b_i$ for all odd i and $b_i \leq a_i$ for all even i . Such a sequence a represents an atom of $K^{\#n}$ iff $a_i = a_{i+1}$ for all odd i ; and a coatom of $K^{\#n}$ iff $a_i = a_{i+1}$ for all even $i < 2^n$, a_1 is an atom of K and a_{2^n} is a coatom of K . Thus the atoms of $K^{\#n}$ can be identified with non-decreasing sequences $a = (a_2 \leq a_4 \leq \dots \leq a_{2^{n-2}} \leq a_{2^n})$ of elements of K , and the coatoms of $K^{\#n}$ with non-decreasing sequences $s = (s_1 \leq s_3 \leq \dots \leq s_{2^{n-1}} \leq s_{2^{n+1}})$ of elements of K , where s_1 is an atom and $s_{2^{n+1}}$ a coatom of K ; in this notation, $a \leq s$ iff $s_1 \leq a_2 \leq s_3 \leq a_4 \leq \dots \leq a_{2^n} \leq s_{2^{n+1}}$. If C is the cover of $|K^{\#n}|$ by the open stars of

vertices of $K^{\#n}$, then the vertices of $N(C)$ correspond to the atoms of $K^{\#n}$, and a set of vertices of $N(C)$ determines a simplex of $N(C)$ iff the corresponding atoms of $K^{\#n}$ all belong to the cone of some coatom of $K^{\#n}$.

Consider the projection $\pi: P_i = [2] + \cdots + [2] \rightarrow [1] + \cdots + [1] \simeq [i]$. By the above, $N(C_n) \subset 2^A$, where A is the set of all non-decreasing sequences $a = (a_2 \leq a_4 \leq \cdots \leq a_{2^n})$ of elements of P_{2^n+1} , and $N(C_n)$ consists of all $S \subset A$ such that there exists a non-decreasing sequence $s = (s_1 \leq s_3 \leq \cdots \leq s_{2^n+1})$ of elements of P_{2^n+1} , where $\pi(s_1) = 1$ and $\pi(s_{2^n+1}) = 2^n + 1$, and $s_1 \leq a_2 \leq s_3 \leq a_4 \leq \cdots \leq a_{2^n} \leq s_{2^n+1}$.

Let $L_k = \{a \in A \mid \pi(a_{2i}) \leq 2i \text{ for all } i \leq k\}$ and $R_k = \{a \in A \mid \pi(a_{2i}) \geq 2i \text{ for all } i \geq 2^{n-1} + 1 - k\}$. Thus $A = L_0 \supset L_1 \supset \cdots \supset L_{2^n-1}$, and $A = R_0 \supset R_1 \supset \cdots \supset R_{2^n-1}$. Note that $L_{2^n-1} \cap R_{2^n-1} = \{a \in A \mid \pi(a_{2i}) = 2i \text{ for all } i\}$, which lies in a single simplex, as witnessed by any sequence s with $\pi(s) = (1 \leq 3 \leq \cdots \leq 2^n + 1)$. For $i = 0, \dots, 2^{n-1}$ let N_i be the full subcomplex of $N(C_n)$ spanned by L_i , and for $i = 2^{n-1} + 1, \dots, 2^n$ let N_i be the full subcomplex of $N(C_n)$ spanned by $L_{2^n-1} \cap R_{i-2^n-1}$. Thus $N_0 = N(C_n)$; on the other hand, since N_{2^n} is a full simplex, $|N_{2^n}|$ is contractible. We shall now construct a deformation retraction of $|N_i|$ onto $|N_{i+1}|$ for each $i = 0, \dots, 2^n - 1$.

We first define a retraction $r_k: L_{k-1} \rightarrow L_k$ by $r_k(a) = a$ if $a \in L_k$, and else by $r_k(a) = b$, where $b_{2i} = a_{2i}$ for $i \neq k$ and $\pi(b_{2k}) = 2k$. (This leaves two possibilities for b_{2k} , among which we choose arbitrarily.) Let $S \in N_{k-1}$ be witnessed by a sequence $s = (s_1 \leq s_3 \leq \cdots \leq s_{2^n+1})$ of elements of P_{2^n+1} , where $\pi(s_1) = 1$ and $\pi(s_{2^n+1}) = 2^n + 1$. If $\pi(s_{2k+1}) \leq 2k$, then all elements of S belongs to L_k , and so $r_k|_S$ is the identity. Else $\pi(s_{2k+1}) \geq 2k + 1$, and since all elements of S belong to L_{k-1} , we may assume that $\pi(s_{2k-1}) \leq 2k - 1$ by modifying s if necessary. Then $r_k(S)$ is a simplex of N_k , and furthermore $S \cup r_k(S)$ is a simplex of N_{k-1} , as witnessed by the same sequence s . Thus r_k extends to a simplicial retraction $R_k: N_{k-1} \rightarrow N_k$, and furthermore we get a simplicial map $H_k: N_{k-1} * N_{k-1} \rightarrow N_{k-1}$ that restricts to the identity on the first factor and to R_k on the second factor. It follows that $|R_k|$ is a deformation retraction.

We next similarly define a retraction $r'_k: R_{k-1} \rightarrow R_k$ by $r'_k(a) = a$ if $a \in R_k$, and else by $r'_k(a) = b$, where $b_{2i} = a_{2i}$ for $i \neq 2^{n-1} + 1 - k$ and $\pi(b_{2^n+2-2k}) = 2^n + 2 - 2k$. (This leaves two possibilities for b_{2^n+2-2k} , among which we choose arbitrarily.) Then $r'_k(L_{2^n-1}) \subset L_{2^n-1}$, and then the preceding argument goes through. This completes the proof that $|N(C_n)|$ deformation retracts onto a contractible subspace, and therefore is itself contractible.

Suppose that $|P|$ satisfies the Hahn property. (The following somewhat technical argument can be somewhat simplified by using Lemma 8.2 below, modulo the proof of that lemma.) Since $|P|$ is uniformly locally contractible, there exists an $\varepsilon > 0$ such that every two ε -close maps into $|P|$ are homotopic with respect to the usual metric on $|P|$. Let D_n be the cover of $|P^{\#n}|$ by the open stars of vertices of $P^{\#n}$. Since $P \simeq VD(D_0)$ is a subposet $N(D_0)$, we have an isometric embedding $|P| \subset |N(D_0)|$ with respect to the usual metrics. We recall that the d_∞ metric on $N(D_0)$ is uniformly equivalent to the usual metric d . Let d' denote the d_∞ metric on $|N(D_0)|$ restricted over $|P|$. Let $\delta > 0$ be such that δ -close points in $|P|$ with respect to d' are ε -close with respect to

d. By our assumption there exists a $\gamma > 0$ such that for every $\beta > 0$, every (β, γ) -continuous map into $(|P|, d')$ is δ -close with respect to d' to a continuous map. Now the $\frac{\gamma}{3}$ -neighborhood U of $|P|$ in $|N(D_0)|$ with respect to the d_∞ metric on $|N(D_0)|$ admits a discontinuous, $(\gamma/3, \gamma)$ -continuous retraction onto $(|P|, d')$. Thus we obtain a continuous map $f: U \rightarrow (|P|, d')$ such that $f|_{|P|}$ is δ -close to the identity. Then by the above $f|_{|P|}$ is homotopic to the identity.

Let n be such that $2^{-n} \leq \gamma/3$. Let $R_n = \lceil VD(D_0)^{\#n} \rceil_{N(D_0)^{\#n}}$. Then $|R_n|$ lies in the 1-neighborhood of $|VD(D_0)^{\#n}|$ with respect to the d_∞ metric on $|N(D_0)^{\#n}|$. Hence the image of $|R_n|$ under the homeomorphism $|N(D_0)^{\#n}| \cong |N(D_0)|$ lies in the 2^{-n} -neighborhood of $|P|$ with respect to the d_∞ metric on $|N(D_0)|$. Thus we obtain a continuous map $g: |R_n| \rightarrow |P|$ whose restriction over $P^{\#n}$ is homotopic to the homeomorphism $|P^{\#n}| \cong |P|$. Next, each D_{i+1} star-refines D_i , so we have the canonical bonding map $\varphi_{D_i}^{D_{i+1}}: N(D_{i+1}) \rightarrow N(D_i)^{\#}$. It is easy to see that the composition $P^{\#(n+1)} \simeq VD(D_{i+1}) \xrightarrow{\varphi} VD(D_i)^{\#n} \simeq P^{\#n}$, where φ is the restriction of $\varphi_{D_i}^{D_{i+1}}$, is the identity map. By iterating we obtain a monotone map $\varphi_n: N(D_n) \rightarrow N(D_0)^{\#n}$ that extends the composition $VD(D_n) \simeq P^{\#n} \simeq VD(D_0)^{\#n}$. Since φ_n is monotone and $N(D_n) = \lceil VD(D_n) \rceil$, the image of φ_n lies in R_n . Thus we obtain a continuous map $h: |N(D_n)| \rightarrow |P|$ whose restriction to $|VD(D_n)|$ is homotopic to the homeomorphism $|P^{\#n}| \cong |P|$. In particular, we obtain a continuous map $k: |N(C_n)| \rightarrow |P_{2^n+1}|$ whose restriction to $|VD(C_n)|$ is homotopic to the homeomorphism $|Q_n| \cong |P_{2^n+1}|$. This yields a continuous retraction of the $(2^n + 1)$ -ball onto the boundary 2^n -sphere, which is a contradiction.

We note that the poset P in Example 7.5 satisfies the following property (Q): There exist essential maps $e_n: S^{2^n} \hookrightarrow |P|$ such that for each $\varepsilon > 0$ there exists an n , a $\delta > 0$ and a discontinuous, (δ, ε) -continuous extension of e_n over B^{2^n+1} . On the other hand, since $|P^b|$ is a uniform ANR, $|P|$ is a non-uniform ANR, and in particular satisfies the non-uniform homotopy extension property. It follows that every metrizable uniform space that is uniformly homotopy equivalent to $|P|$ satisfies (Q) as well. In particular, using that $|P|$ is uniformly locally contractible, we get the following

Theorem 7.6. *There exists a countable poset whose geometric realization is not uniformly homotopy equivalent to a uniform ANR, nor even to a metrizable uniform space satisfying the Hahn property.*

The remainder of this subsection is not used elsewhere in this paper, and could be of interest primarily to the reader who is looking for a class of posets larger than conditionally complete posets whose geometric realizations are uniform ANRs.

7.7. Hereditariness. We call a cover D of a metric space M *hereditarily uniform*, if there exists a $\lambda > 0$ such that each $E \subset D$ is a uniform cover of $\bigcup E$ with Lebesgue number λ . Any such λ is a *hereditary Lebesgue number* of the hereditarily uniform cover.

We say that a cover C of a set S *hereditarily star-refines* a cover D of S if for every $E \subset D$, the cover $C \cap (\bigcup E)$ of the subset $\bigcup E \subset X$ star-refines the cover E of $\bigcup E$.

If D is a cover of M by sets of diameters $< \varepsilon$, and C is a hereditarily uniform cover of M with a hereditary Lebesgue number 2ε , then clearly D hereditarily star-refines C .

Lemma 7.8. *Let C and D be covers of a set S . If C hereditarily star-refines D , then φ_D^C sends $N(C)$ into $VD(D)^\#$.*

Proof. The hypothesis implies that for every $x \in C$, every $T \subset \text{st}(x, C)$ such that $x \in T$ satisfies the following property (*): if T lies in $\bigcup E$ for some $E \subset D$, then it lies in some element of E . In particular, (*) is satisfied by any T of the form $\bigcap \sigma$ or $\bigcup \sigma$, where $\sigma \in N(C)$. On the other hand, every element of $N(D)$ of the form $\Delta_D(T)$ where T satisfies (*) clearly belongs to $VD(D)$. $\square$

7.9. Construction of hereditary uniform covers. For a finite-dimensional atomic poset P with atom set Λ , it is easy to construct a hereditarily uniform cover of $|P|$, namely the cover by the sets $U_\lambda = \bigcup_{\sigma \geq \lambda} |H_\sigma|$ composed of the barycentric handles $H_\sigma = \lceil \langle \sigma \rangle^* \rceil \subset (P^b)^*$, where $\sigma \in P$. The hereditary is due to the fact that $|H_\sigma|$ and $|H_\tau|$ are uniformly disjoint when σ and τ are incomparable and P is finite-dimensional. This argument does not apply to canonical handles $h_\sigma = \lceil [\sigma, \sigma]^* \rceil \subset (P^\#)^*$ because $h_\sigma \cap h_\tau$ can be nonempty when σ and τ are incomparable.

Clearly, the preimage of a hereditarily uniform cover under a uniformly continuous map of metrizable uniform spaces is hereditarily uniform. Hence by using Lemma 6.7, we infer that every uniform cover of a residually finite-dimensional metrizable uniform space admits a hereditarily uniform refinement (in fact, one of a finite multiplicity). We conjecture that the hypothesis of residual finite-dimensionality cannot be dropped here.

7.10. Weak hereditary. We call a cover D of a metric space M *weakly hereditarily uniform*, if there exists a $\lambda > 0$ such that for every $F \subset D$ satisfying $\bigcap F \subset \bigcup (D \setminus F)$, the cover $(D \setminus F) \cap (\bigcap F)$ of the subset $\bigcap F \subset X$ is a uniform cover of $\bigcap F$ with Lebesgue number λ . Any such λ is called a *weak hereditary Lebesgue number* of the weakly hereditarily uniform cover. A hereditarily uniform cover is weakly hereditarily uniform by considering $E = D \setminus F$; and a weakly hereditarily uniform cover is uniform by considering $F = \emptyset$ (in which case $\bigcap F = M$).

We say that a cover C of a set S *weakly hereditarily star-refines* a cover D of S if for every $F \subset D$ satisfying $\bigcap F \subset \bigcup (D \setminus F)$, the cover $C \cap (\bigcap F)$ of the subset $\bigcap F \subset X$ star-refines the cover $(D \setminus F) \cap (\bigcap F)$ of $\bigcap F$. Similarly to the above,

$$\text{hereditary star-refinement} \Rightarrow \text{weak hereditary star-refinement} \Rightarrow \text{star-refinement}.$$

It is easy to see that if D is a cover of M by sets of diameters $< \varepsilon$, and C is a weakly hereditarily uniform cover of M with a weak hereditary Lebesgue number 2ε , then D weakly hereditarily star-refines C .

Beware that the preimage of a weakly hereditarily uniform cover under a uniformly continuous map f of metrizable uniform spaces need not be weakly hereditarily uniform, because $\bigcap F \not\subset \bigcup (D \setminus F)$ does not imply $f^{-1}(\bigcap F) \not\subset f^{-1}(\bigcup (D \setminus F))$.

The proof of Lemma 7.8 works to establish

Lemma 7.11. *Let C and D be covers of a set S . If C weakly hereditarily star-refines D , then φ_D^C sends $N(C)$ into $VD(D)^\#$.*

8. APPROXIMATION OF SPACES

Theorem 8.1. *Every separable metrizable complete uniform space is the limit of a convergent inverse sequence of geometric realizations of simplicial complexes and uniformly continuous maps.*

Theorem 8.1 follows from [13; Theorem 2.42] along with the following

Lemma 8.2. *Let $\{C_n\}$ be a basis of the uniformity of a metrizable complete uniform space X , where each uniform cover C_n is countable and point-finite. Let $N_i = |N(C_i)|$, and let $p_i: N_{i+1} \rightarrow N_i$ be the composition $|N(C_{i+1})| \xrightarrow{|\varphi_{C_i}^{C_{i+1}}|} |N(C_i)^\#| \xrightarrow{h} |N(C_i)|$, where $\varphi_{C_i}^{C_{i+1}}$ is the canonical bonding map, and h is the uniform homeomorphism. Then the inverse sequence $\dots \xrightarrow{p_1} N_1 \xrightarrow{p_0} N_0$ is convergent, and its limit L is uniformly homeomorphic to X .*

Though our notion of geometric realization and our bonding maps are quite different from those used by Isbell, our proof of Lemma 8.2 is modelled on Isbell's proof of what amounts to its residually finite-dimensional case [10; Lemma V.33].⁷

Proof. Let $s_i(x)$ denote the simplex $[\Delta_{C_i}(x)]$. Then each $f_i := \varphi_{C_i}^{C_{i+1}}$ sends $s_{i+1}(x)$ into $s_i(x)^\#$ for each $x \in X$. Hence each p_i sends $|s_{i+1}(x)|$ into $|s_i(x)|$. Since f_i is monotone, $|f_i|$ is 1-Lipschitz with respect to the d_∞ metrics, and therefore p_i is $\frac{1}{2}$ -Lipschitz with respect to the d_∞ metrics. Since the diameter of $|s_i(x)|$ is bounded above by 2 in the d_∞ metric, the diameter of $p_i^{i+n}(|s_{i+n}(x)|)$ is bounded above by 2^{1-n} . Since each $|s_i(x)|$ is compact, their inverse limit (with the restrictions of p_i as the bonding maps) is nonempty, and by the above it has zero diameter. Thus it is a single point $\lambda(x) \in L$.

Each N_i is the union of the $|s_i(x)|$ over all $x \in X$, and every $|s_i(x)|$ contains $p_i^\infty(\lambda(x))$. Hence every point of $p_i^{i+n}(N_{i+n})$ is 2^{1-n} -close to a point of $p_i^\infty(L)$. Thus the inverse sequence is convergent (see [13; Lemma 5.6(c)]).

To see that $\lambda: X \rightarrow L$ is uniformly continuous, it suffices to show that every its coordinate $\lambda_i: X \xrightarrow{\lambda} L \xrightarrow{p_i^\infty} N_i$ is uniformly continuous. Indeed, for each $x \in X$ and each $j \geq i$ we have $\lambda_j(x) \in |s_j(x)| = |[\Delta_{C_j}(x)]|$. For each $V \in C_j$, every $x \in V$ satisfies $V \in \Delta_{C_j}(x)$. Hence $\lambda_j(V) \subset |\text{st}(\{V\}, N(C_j))|$. The diameter of $|\text{st}(\{V\}, N(C_j))|$ is bounded above by 4, hence its image under p_i^j has diameter at most $2^{2-(j-i)}$. Thus for each $\varepsilon > 0$ there exists a $j \geq i$ such that $\lambda_i(C_j) = p_i^j \lambda_j(C_j)$ refines the cover of N_i by ε -balls. Since $\{C_j\}$ is a fundamental sequence of covers of X , we infer that λ_i is uniformly continuous.

⁷There is a minor error in the proof of step (2) in [10; Lemma V.33], as the Cauchy filter base considered there might consist entirely of the empty sets. This can be remedied as shown in the last paragraph of our proof.

Next, given $x, y \in X$ at a distance $\varepsilon > 0$, there exists an $n = n(\varepsilon)$ such that any two elements of C_n containing x and y are disjoint. Then λ_n sends x and y into disjoint closed simplices $|s_n(x)|$ and $|s_n(y)|$ of N_n . It follows that λ is injective and, using the uniform continuity of each p_n^∞ , that λ^{-1} is uniformly continuous.

Finally, if $(q_i) \in L$ is a thread of $q_i \in N_i$, and σ_n is the minimal simplex of $N(C_n)$ such that $q_n \in |\sigma_n|$, then $f_n(\sigma_{n+1}) \subset \sigma_n^\#$, moreover, σ_n is the minimal simplex of $N(C_n)$ satisfying the latter property. Hence $\sigma_n = \Delta_{C_n}(\bigcap \sigma_{n+1})$; in particular, $\bigcap \sigma_{n+1} \subset \bigcap \sigma_n$. Let S_n be the closure of $\bigcap \sigma_n$. Then S_n lies in the closure of an element of C_n . Since $\{C_i\}$ is a basis of the uniformity of X , for each $\varepsilon > 0$ there exists an n such that every element of C_n is of diameter at most ε . It follows that the inverse sequence $\cdots \subset S_1 \subset S_0$ is Cauchy (see [13; Lemma 5.6(d)]). Since X is complete, so are the S_i 's, hence $\cdots \subset S_1 \subset S_0$ is convergent (see [13; Lemma 5.6(b)]) and therefore $\bigcap S_i$ is nonempty (see [13; Lemma 5.6(f)]). Since the diameters of S_i tend to zero, $\bigcap S_i$ must be a single point q . Now q lies in the closure of $\bigcap \sigma_n$, and each $q' \in \bigcap \sigma_n$ satisfies $\sigma_n \subset s_n(q')$ and $\lambda_n(q') \in |s_n(q')|$. Since λ_n is continuous, $\lambda_n(q)$ lies in the closed subset $|\lceil \sigma_n \rceil|$ of $|N(C_n)|$. Hence $s_n(q) \subset \lceil \sigma_n \rceil$, or equivalently $\sigma_n \subset \lceil s_n(q) \rceil$. Since $\lambda(q)$ is also the inverse limit of the simplicial neighborhoods $|\lceil s_n(q) \rceil|$, we conclude that $\lambda(q) = (q_i)$. Thus λ is surjective. $\square$

Remark 8.3. Given simplicial complexes K_i and monotone maps $f_i: K_{i+1} \rightarrow K_i^\#$, it is easy to see (by analyzing the definition of a countable product of uniform spaces, see [13]) that a base of uniformity of $\varprojlim |K_i|$ is given by the preimages of the covers of $|K_i|$'s by the open stars of vertices. It follows, for instance, that $\text{id}: \varprojlim |K_i| \rightarrow \varprojlim |K_i^{\#n}|$ is 2^n -Lipschitz. The best that can be said of the inverse is that it is 1-Lipschitz.

Remark 8.4. It should be noted that the theory of inverse limits traces back ultimately to the work of Alexandroff [1], who had obtained a version of Lemma 8.2 with mere refinement instead of the star-refinement in the hypothesis, and with every thread, as a point of the inverse limit in the conclusion, replaced by the set of the simplicial neighborhoods of its elements.

Similarly to [9; proof of 7.2(ii)], one deduces from Theorem 8.1 the following

Corollary 8.5. *If $\{C_\alpha\}$ is a basis of uniform covers of a separable complete uniform space X , where each C_α is countable and point-finite, then X is a (non-sequential) inverse limit of inverse limits of uniformly continuous maps between $|N(C_\alpha)|$.*

Theorem 8.6. *Every separable complete uniform space is the limit of an inverse spectrum of uniform polyhedra and uniformly continuous maps.*

The following argument is an elaboration on [9; 7.2(i)]. It shows that the uniform polyhedra can be chosen to be geometric realizations of cubical complexes, and more specifically cubohedra in the sense of [13].

Proof. It is well-known that every uniform space X embeds in an (uncountable) product of complete metric spaces M_i [10; II.14, II.15]. If X is separable, we may assume that

so is each M_i , by considering the closures of the images of X in the M_i 's. Then by Theorem 8.1, each M_i in turn embeds in a product of uniform polyhedra. Thus X can be identified with a subspace of a product of uniform polyhedra $|K_j|$.

Since $\prod |K_j|$ is the inverse limit of finite subproducts, and X is closed in $\prod |K_j|$, it is the inverse limit of its images in the finite subproducts, cf. [10; IV.34]. Each finite subproduct is a uniform polyhedron $|K_{j_1} \times \dots \times K_{j_r}|$. Let $P_{n;j_1\dots j_r}$ be the minimal subcomplex of $(K_{j_1} \times \dots \times K_{j_r})^{\#n}$ such that $|P_{n;j_1\dots j_r}|$ contains the image of X . Then it is shown similarly to [5; proof of Theorem 10.1] that X is the inverse limit of all the $|P_{n;j_1\dots j_r}|$. $\square$

Theorem 8.7. *A separable metric space is a uniform ANR if and only if it is uniformly ε -homotopy dominated by the geometric realization of a simplicial complex, for each $\varepsilon > 0$.*

The “if” direction follows from [13; Corollary 4.23]. The following proof of the “only if” direction is based on Theorem 8.1; the reader who feels that this is an overkill can get a more elementary argument by combining the first sentence of this proof with the proof of Theorem 8.8 below.

Proof, “only if”. By [13; Theorem 4.9], the given uniform ANR is uniformly ε -homotopy equivalent to its completion X , which is still a uniform ANR, for each $\varepsilon > 0$. By Theorem 8.1, X is the limit of a convergent inverse sequence of geometric realizations P_i of simplicial complexes, and uniformly continuous bonding maps p_i . By [13; Corollary 5.20] there exists a k and a uniformly continuous retraction $r_{[k,\infty]}: P_{[k,\infty]} \rightarrow X$. For each $l \geq k$ let r_l and $r_{[l,\infty]}$ denote the restrictions of $r_{[k,\infty]}$ over P_l and over $P_{[l,\infty]}$, respectively. Let $p_{[k,\infty]}^\infty: X \times I \rightarrow P_{[k,\infty]}$ be obtained by combining the maps $p_i^\infty: X \rightarrow P_i$. Then for each $l \geq k$, the composition $r_{[l,\infty]} p_{[l,\infty]}^\infty: X \times I \rightarrow X$ is a uniformly continuous homotopy between $X \xrightarrow{p_l^\infty} P_l \xrightarrow{r_l} X$ and id_X . Moreover, for each $\varepsilon > 0$ there exists a k such that $r_{[l,\infty]} p_{[l,\infty]}^\infty$ is an ε -homotopy. Thus X is uniformly ε -homotopy dominated by P_l . $\square$

Theorem 8.8. *Every separable ANRU X is uniformly C -homotopy dominated by the geometric realization of a simplicial complex, for each uniform cover C of X .*

The following proof elaborates on [9; proof of 7.3].

Proof. Since X is an ANRU it is uniformly locally contractible, i.e. for every uniform cover C of X there exists a uniform cover C_1 of X such that for every uniform space Y , every two C_1 -close maps $Y \rightarrow X$ are uniformly C -homotopic [9; proof of 4.2]. Let C_2 be a uniform star-refinement of C_1 . Since X satisfies the Hahn property [9; 4.1], there exists a uniform cover C_3 of X such that for any uniform space Y , any map $\varphi: Y \rightarrow X$ such that $\varphi^{-1}(C_3)$ is uniform is C_2 -close to a uniformly continuous map. Let C_4 be a uniform strong star-refinement of C_3 .

By [10; Theorem I.14] there exists a uniformly continuous map f from X onto a metric space M such that C_4 is refined by $f^{-1}(D)$, where D is the cover of M by all sets of diameter < 1 . Since X is separable and f is surjective, M is separable. Then D is

refined by a point-finite countable uniform cover E (see [13; Theorem 2.42]). Without loss of generality E is open. Then by Lemma 6.7 there exists a uniformly continuous map $g: M \rightarrow |N(E)|$ such that $E = f^{-1}(F)$, where F is the cover of $|N(E)|$ by the open stars of vertices. Let us define a discontinuous map $\varphi: |N(E)| \rightarrow X$ by sending an $x \in |N(E)|$ to any point in $f^{-1}(g^{-1}(U))$, where U is any element of F containing x . Note that $g^{-1}(U)$ is nonempty, being an element of E ; hence $f^{-1}(g^{-1}(U))$ is nonempty, due to the surjectivity of f . Since $f^{-1}(g^{-1}(U))$ is contained in some $V_U \in C_4$, we get that $\varphi(F)$ refines C_3 and the composition $X \xrightarrow{f} M \xrightarrow{g} |N(E)| \xrightarrow{\varphi} X$ is C_4 -close to the identity. Then there exists a uniformly continuous map $\psi: |N(E)| \rightarrow X$ that is C_2 -close to φ . Hence ψ is C_1 -close to the identity, and therefore C -homotopic to it. $\square$

The following improves on [13; Theorems 4.20 and 4.29].

Theorem 8.9. *A separable uniform space X satisfies the Hahn property if and only if for each uniform cover C of X there exists a simplicial complex K and uniformly continuous maps $X \xrightarrow{f} |K| \xrightarrow{g} X$ whose composition is C -close to id_X .*

Proof. The “only if” assertion can be proved similarly to Theorem 8.8 (or, in the metrizable case, to Theorem 8.7). The “if” assertion is easy (cf. [13]). $\square$

Theorem 8.10. *If X is a uniform ANR, then $X \times \mathbb{R}$ is uniformly homotopy equivalent to a uniform polyhedron.*

Proof. By Theorem 8.7 we are given uniformly continuous maps $d: |K| \rightarrow X$ and $u: X \rightarrow |K|$, where K is a simplicial complex, such that the composition $X \xrightarrow{u} |K| \xrightarrow{d} X$ is uniformly homotopic to the identity by an homotopy $h: X \times I \rightarrow X$. We now perform a uniform version of Mather’s trick (see [6]): $X \times \mathbb{R}$ is uniformly homotopy equivalent to the double mapping telescope of

$$\cdots \rightarrow X \xrightarrow{du} X \xrightarrow{du} X \rightarrow \cdots,$$

which is in turn uniformly homotopy equivalent to the double mapping telescope of

$$\cdots \rightarrow X \xrightarrow{d} |K| \xrightarrow{u} X \xrightarrow{d} |K| \xrightarrow{u} X \rightarrow \cdots,$$

which is in turn uniformly homotopy equivalent to the double mapping telescope of

$$\cdots \rightarrow |K| \xrightarrow{ud} |K| \xrightarrow{ud} |K| \rightarrow \cdots.$$

Since $|K|$ is a uniform ANR, by [13; Lemma 4.16] it is uniformly locally contractible. Then by Theorem 7.4, ud is uniformly homotopic to the composition $|K| \xrightarrow{h_m^{-1}} |K^{\#m}| \xrightarrow{|f|} |K^{\#n}| \xrightarrow{h_n} |K|$ for some monotone map $f: K^{\#m} \rightarrow K^{\#n}$, where $m \geq n$ for the sake of definiteness. On the other hand, we have the monotone map $\#(m-n): K^{\#m} \rightarrow K^{\#n}$, whose geometric realization is uniformly homotopic to the uniform homeomorphism $h_m: |K^{\#m}| \rightarrow |K^{\#n}|$ (see Lemma 4.1). Thus $X \times \mathbb{R}$ is uniformly homotopy equivalent to the geometric realization of the double mapping telescope of

$$\cdots \xleftarrow{\#(m-n)} K^{\#n} \xrightarrow{f} K^{\#m} \xleftarrow{\#(m-n)} K^{\#n} \xrightarrow{f} \cdots$$

By Theorem 4.12, the latter is uniformly homotopy equivalent to the geometric realization of the thickened double mapping telescope, which is a uniform polyhedron (using [12; Lemma 2.21]). $\square$

Remark 8.11. Similar arguments (with double mapping telescopes not of individual nerves but of their mapping telescopes) also show that if X is a complete uniform ANR, then $X \times \mathbb{R}$ is the limit of a convergent inverse sequence $\dots \xrightarrow{q_1} Q_1 \xrightarrow{q_0} Q_0$ of geometric realizations of countable preposets and uniformly continuous maps such that each q_i is a uniform homotopy equivalence.

Theorem 8.12. *Every complete uniform ANR is the limit of a convergent inverse sequence $\dots \xrightarrow{q_1} Q_1 \xrightarrow{q_0} Q_0$ of uniform polyhedra and uniformly continuous maps such that each q_i is (non-uniformly) a homotopy equivalence.*

Proof. Let X be the given complete uniform ANR. By Theorem 8.1, X is the limit of a convergent inverse sequence $\dots \xrightarrow{p_1} P_1 \xrightarrow{p_0} P_0$ of uniform polyhedra and uniformly continuous maps. Suppose that we have constructed a finite sequence $n_0, n_1, n_2, \dots, n_k$ and a finite chain of uniform polyhedra and uniformly continuous maps $P_{n_k} \subset Q_k \xrightarrow{q_{n-1}} \dots \xrightarrow{q_1} P_{n_1} \subset Q_1 \xrightarrow{q_0} P_{n_0} \subset Q_0$ such that the composition $X \xrightarrow{p_{n_i}^\infty} P_{n_i} \subset Q_i$ is a homotopy equivalence for each $i \leq k$.

Since P_{n_k} is uniformly locally contractible, there exists an $\varepsilon > 0$ such that every two ε -close uniformly continuous maps $Y \rightarrow P_{n_k}$ are uniformly homotopic. Let δ be such that p_i^∞ sends δ -close points into $\varepsilon/2$ -close points. Since the inverse sequence is convergent and X satisfies the Hahn property (see [13; Lemma 4.19(a)]), there exists an $m \geq n$ and a map $r: P_m \rightarrow X$ such that the composition $X \xrightarrow{p_m^\infty} P_m \xrightarrow{r} X$ is δ -close to the identity. Then the composition $X \xrightarrow{p_m^\infty} P_m \xrightarrow{r} X \xrightarrow{p_{n_k}^\infty} P_{n_k}$ is $\varepsilon/2$ -close to $p_{n_k}^\infty$. Since the inverse sequence is convergent, there exists an $l \geq m$ such that the composition $P^l \xrightarrow{p_m^l} P_m \xrightarrow{r} X \xrightarrow{p_{n_k}^\infty} P_{n_k}$ is ε -close to $p_{n_k}^l$. Let d be the composition $P_l \xrightarrow{p_m^l} P_m \xrightarrow{r} X$, let $u = p_l^\infty$, and let f be the composition $P_l \xrightarrow{d} X \xrightarrow{u} P_l$. Then the composition $P_l \xrightarrow{f} P_l \xrightarrow{p_{n_k}^l} P_{n_k}$ is uniformly homotopic to $p_{n_k}^l$. This yields a uniformly continuous map $F: MC(f) \rightarrow P_{n_k}$ that restricts to $p_{n_k}^l$ on each of the two copies of P_l in $MC(f)$. Let $n_{k+1} = l$. Applying to the maps $X \xrightarrow{u} P_l \xrightarrow{d} X$ the construction in the proof of Theorem 8.10, we obtain the double mapping telescope Q_l homotopy equivalent to X , via the composition $X \xrightarrow{u} P_l \subset Q_{k+1}$. The partial map $Q_{k+1} \supset P_l \xrightarrow{p_{n_k}^l} P_{n_k}$ now extends (using F) to a total uniformly continuous map $q_{k+1}: Q_{k+1} \rightarrow P_{n_k}$, and we are done with the inductive step.

The assertion now follows using that inverse limit is unchanged upon passage to an infinite subsequence. $\square$

REFERENCES

- [1] P. S. Alexandroff, *Simpliziale Approximationen in der allgemeinen Topologie*. Math. Ann. **96** (1926), 489–511; *Bemerkung zu die Arbeit*, Math. Ann. **101** (1929), 452–456. Translated into Russian in Alexandroff’s selected works edition (vol. 1 “Theory of functions of real variable and theory of topological spaces”, publ. 1978). [↑6.1, 8.4](#)
- [2] M. R. Bridson and A. Haefliger, *Metric Spaces of Non-Positive Curvature*, Springer, Berlin, 1999. [↑3](#)
- [3] D. Burago, Yu. Burago, and S. Ivanov, *A Course in Metric Geometry*, Grad. Stud. in Math. vol. 33, Amer. Math. Soc. Providence, RI, 2001; Russian transl. in Institut komputernyh issl. Moscow/Izhevsk, 2004. [↑3](#)
- [4] H. H. Corson and J. R. Isbell, *Some properties of strong uniformities*, Quart. J. Math. Oxford Ser. (2) **11** (1960), 17–33. [↑1.A, 1.B](#)
- [5] S. Eilenberg and N. Steenrod, *Foundations of Algebraic Topology*, Princeton Univ. Press, Princeton, NJ, 1952. [↑8](#)
- [6] S. Ferry and A. Ranicki, *A survey of Wall’s finiteness obstruction*, Surveys on Surgery Theory, vol. 2 (S. Cappell, A. Ranicki, and J. Rosenberg, eds.) Princeton Univ. Press, 2000. [Ranicki’s webpage](#), pp. 63–80. [↑8](#)
- [7] H. Freudenthal, *Entwicklungen von Räumen und ihren Gruppen*, Compositio Math. **4** (1937), 145–234. [↑1.A](#)
- [8] J. R. Isbell, *On finite-dimensional uniform spaces*, Pacific J. Math. **9** (1959), 107–121. [↑1.1, 1.A, 1.B, 4](#)
- [9] ———, *Uniform neighborhood retracts*, Pacific J. Math. **11** (1961), 609–648. [↑1.A, 1.B, 8, 8, 8](#)
- [10] ———, *Uniform Spaces*, Amer. Math. Soc. Providence, RI, 1964. [↑1.A, 1.B, 4, 6, 8, 7, 8, 8](#)
- [11] ———, *Irreducible polyhedral expansions*, Nederl. Akad. Wetensch. Proc. Ser. A 64 = Indag. Math. **23** (1961), 242–248. [↑1.A](#)
- [12] S. A. Melikhov, *Combinatorics of combinatorial topology*. [arXiv:1208.????](#) [↑1, 1.1, 1.A, 1.A, 1.C, 2, 2.1, 2.2, 2.3, 2.9, 2, 3, 3.5, 3.6, 4, 4, 4, 6, 6.19, 8](#)
- [13] ———, *Metrisable uniform spaces*. [arXiv:1106.3249](#). [↑1, 1.A, 1.A, 1.A, 1.B, 1.C, 2, 3, 3.4, 4.6, 4, 4, 4.9, 4, 6, 7, 8, 8, 8.3, 8, 8, 8, 8, 8, 8](#)
- [14] M. A. Shtan’ko and M. I. Shtogrin, *On the embedding of cubic manifolds and complexes in a cubic lattice*, Uspekhi Mat. Nauk **47:1** (1992), 219–220; English transl., Russ. Math. Surv. **47:1** (1992), 267–268. [↑1.A, 2](#)

STEKLOV MATHEMATICAL INSTITUTE OF THE RUSSIAN ACADEMY OF SCIENCES, UL. GUBKINA 8, MOSCOW, 119991 RUSSIA

E-mail address: melikhov@mi.ras.ru

DELAUNAY LABORATORY OF DISCRETE AND COMPUTATIONAL GEOMETRY, YAROSLAVL’ STATE UNIVERSITY, 14 SOVETSKAYA ST., YAROSLAVL’, 150000 RUSSIA